

Asymmetric and Moving-Frame Approaches to MHD Equations

Bintao Cao*

Abstract

The magnetohydrodynamic (MHD) equations of incompressible viscous fluids with finite electrical conductivity describe the motion of viscous electrically conducting fluids in a magnetic field. In this paper, we find twelve families of solutions of these equations by Xu's asymmetric and moving frame methods. A family of singular solutions may reflect basic characteristics of vortices. The other solutions are globally analytic with respect to the spacial variables. In particular, Bernoulli equation and Wronskian determinants play important roles in our approaches. Our solutions may also help engineers to develop more effective algorithms to find physical numeric solutions to practical models.

Keywords: MHD equations; asymmetric condition; moving frame; Bernoulli equation.

AMS Subject Classification (2000): 35Q51, 35C10, 35C15.

1 Introduction

In physics, magnetohydrodynamic (MHD) studies the motion of electrically conducting fluids in a magnetic field. Examples are astrophysical plasma, solar atmosphere, magnetosphere, thermonuclear fusion and liquid metal etc. Their motion described by the MHD equations (see [1] for the details), which coupled the Navier-Stokes equations and the Maxwell's equations.

There are many different approaches to the MHD equations. The wellposedness of these equations has been well studied (e.g., cf. [2, 3, 4, 5, 6, 7]). Numerical methods for compressible case were developed in [8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18], etc. One can also find similar results for the incompressible case in [1, 19, 20, 21, 22, 23]. The Lagrange invariance of a generalized ion magnetic helicity was established for Hall MHD in [24]. Vodák [25] proved the existence of a weak, local in time solution to the compressible case on the whole space under special assumptions on pressure and entropy. For the historic introduction of the MHD equations, we refer the Section 1 of [26].

In this paper, we investigate an incompressible case, that is, the MHD equations of incompressible viscous fluids with finite electrical conductivity (e.g., cf. [27]):

$$\nabla \cdot \mathbf{v} = 0, \quad (1.1)$$

$$\mathbf{v}_t + (\mathbf{v} \cdot \nabla) \mathbf{v} - \mu_0 (\mathbf{H} \times \text{rot} \mathbf{H}) + \frac{1}{\rho} \nabla p = \nu \Delta \mathbf{v}, \quad (1.2)$$

$$\nabla \cdot \mathbf{H} = 0, \quad (1.3)$$

$$\mathbf{H}_t = \text{rot}(\mathbf{v} \times \mathbf{H}) + \eta \Delta \mathbf{H}. \quad (1.4)$$

Here $\mathbf{v} = (u, v, w)^t$ is the velocity vector, $\mathbf{H} = (H^1, H^2, H^3)^t$ is the magnetic field vector, ν is the viscosity coefficient, μ_0 is the magnetic permeability and η is the magnetic viscosity. The superscript “ t ” denotes the transpose of a matrix or a vector.

*Institute of Mathematics, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, P. R. China, caobintao@amss.ac.cn

Nucci [28] found the Lie point symmetries of these equations. Popovych [29, 30, 31] reduced (1.1)-(1.4) to ordinary differential equations and to partial differential equations in two and in three independent variables by means of Lie symmetries. Then he constructed some classes of similarity solutions.

In order to solve the three-dimensional Navier-Stokes equations, Xu [32] introduced a method of imposing asymmetric conditions on the velocity vector with respect to independent variables and a method of moving frame which reflect the rotation nature of flow. By these methods, he found seven families of non-steady rotating asymmetric solutions with multiple parameter functions. In particular, a family of singular solutions may reflect important features of vortices. The other solutions are globally analytic with respect to spacial variables. The goal of this work is to find similar solutions for the MHD equations (1.1)-(1.4).

Note that when $\mathbf{H} = \vec{0}$, the MHD equations (1.1)-(1.4) become the three-dimensional Navier-Stokes equations. The factor of magnetic field makes the MHD equations much more complicated than the Navier-Stokes equations. For instance, Xu [32] reduced the Navier-Stokes equations under certain conditions to a linear differential equation, and we can only reduce the MHD equations to a system of linear differential equations. In particular, we have to use Wronskian determinants to solve such systems. Moreover, the magnetic field also naturally leads the appearance of Bernoulli equation in our reductions.

Since the system of MHD equations is a coupled system of Navier-Stokes equations and the Maxwell's equations, Xu's methods and exact solutions of the Navier-Stokes equations do give us hints of how to solve the MHD equations exactly. We find twelve families of solutions of the equations with multiple parameter functions. One of them is a family of singular solutions that may reflect important characteristics of vortices. The other solutions are globally analytic with respect to the spacial variables, related to polynomials, exponential functions or trigonometric functions. Our solutions may help mathematicians to develop the method of proving the well known regularity of the equations. They may also help engineers to develop more effective algorithms to find physical numeric solutions to practical models. Below we give more technical details.

For convenience, we always assume that all the involved partial derivatives of related functions always exist and we can change the order of taking partial derivatives. We also use prime ' to denote the derivative of any one-variable function.

According to [28], the MHD equations are invariant under the following transformations:

1. Time translation:

$$T_1 : \quad t \mapsto t + a, \quad (1.5)$$

where a is an arbitrary real constant.

2. Scale transformation:

$$T_2 : \quad (t, \mathbf{x}, \mathbf{v}, \mathbf{H}, p) \mapsto (\lambda^2 t, \lambda \mathbf{x}, \lambda \mathbf{v}, \lambda^{-1} \mathbf{H}, \lambda^{-2} p), \quad (1.6)$$

where λ is any nonzero real constant and $\mathbf{x} = (x, y, z)^t$.

3. Rotations:

$$T_3 : \quad (t, \mathbf{x}, \mathbf{v}, \mathbf{H}, p) \mapsto (t, R\mathbf{x}, R\mathbf{v}, R\mathbf{H}, p) \quad (1.7)$$

for any $R \in SO(3)$.

4. Moving coordinates:

$$T_4 : \quad (t, \mathbf{x}, \mathbf{v}, \mathbf{H}, p) \mapsto (t, \mathbf{x} + \mathbf{F}(t), \mathbf{v} + \mathbf{F}'(t), \mathbf{H}, p - \rho \mathbf{x} \cdot \mathbf{F}''), \quad (1.8)$$

where $\mathbf{F}(t) = (f(t), g(t), h(t))^t$ is an arbitrary vector valued functions.

5. Pressure changes:

$$T_5 : \quad (t, \mathbf{x}, \mathbf{v}, \mathbf{H}, p) \mapsto (t, \mathbf{x}, \mathbf{v}, \mathbf{H}, p + \theta(t)) \quad (1.9)$$

for any function θ of t .

The above transformations transform a solution of the MHD equations into other solutions. So we only need to find exact solutions modulo these transformations.

Assuming

$$\begin{cases} u = \frac{\alpha}{2}x + y\phi(t, x^2 + y^2), \\ v = \frac{\alpha}{2}y - x\phi(t, x^2 + y^2), \\ w = -\alpha z + \psi(t, x^2 + y^2), \end{cases} \quad \begin{cases} H^1 = \frac{\beta}{2}x + y\xi(t, x^2 + y^2), \\ H^2 = \frac{\beta}{2}y - x\xi(t, x^2 + y^2), \\ H^3 = -\beta z + \sigma(t, x^2 + y^2) \end{cases} \quad (1.10)$$

for some functions α and β of t , and two-variable functions ϕ, ψ, ξ, σ , we obtain the following solution:

$$\begin{cases} u = \frac{\alpha}{2}x + y \sum_{m=0}^n a_m(t)(x^2 + y^2)^m, \\ v = \frac{\alpha}{2}y - x \sum_{m=0}^n a_m(t)(x^2 + y^2)^m, \\ w = -\alpha z + \sum_{m=0}^n g_m(t)(x^2 + y^2)^m, \end{cases} \quad \begin{cases} H^1 = \frac{b}{2}x + y \sum_{m=0}^n c_m(t)(x^2 + y^2)^m, \\ H^2 = \frac{b}{2}y - x \sum_{m=0}^n c_m(t)(x^2 + y^2)^m, \\ H^3 = -bz + \sum_{m=0}^n f_m(t)(x^2 + y^2)^m. \end{cases} \quad (1.11)$$

and p is given by (3.86), where n is any given positive integer,

$$\alpha = \frac{k + 2k^2le^{-kt}}{-1 + 2kle^{-kt}} \quad (1.12)$$

with constants k and l is obtained by solving a related Bernoulli equation, b is an arbitrary real constant, and $a_m(t), c_m(t), f_m(t), g_m(t)$ are determined via various Wronskian determinants (cf. (3.76)-(3.85)). This is an example of asymmetric approach.

By Xu's moving-frame approach, we obtain the following solution of the MHD equations (1.1)-(1.4):

$$\begin{cases} u = -(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'})x - \alpha'y - \frac{6\nu \cos \alpha}{x \cos \alpha + y \sin \alpha} + 6\nu \sin \alpha \frac{-x \sin \alpha + y \cos \alpha}{(x \cos \alpha + y \sin \alpha)^2}, \\ v = \alpha'x - (\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'})y - \frac{6\nu \sin \alpha}{x \cos \alpha + y \sin \alpha} - 6\nu \cos \alpha \frac{-x \sin \alpha + y \cos \alpha}{(x \cos \alpha + y \sin \alpha)^2}, \\ w = 2(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'})z, \end{cases} \quad (1.13)$$

$$\begin{cases} H^1 = -[(a \cos \alpha - b \sin \alpha)x + (-a \sin \alpha + b \cos \alpha)y] \sin \alpha, \\ H^2 = [(a \cos \alpha - b \sin \alpha)x + (a \sin \alpha + b \cos \alpha)y] \cos \alpha, \\ H^3 = -bz \end{cases} \quad (1.14)$$

and

$$\begin{aligned} p = & -\rho \left(\left(\left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)' + \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)^2 - (\alpha')^2 - \mu_0 a^2 \right) \frac{(x \cos \alpha + y \sin \alpha)^2}{2} \right. \\ & + \left(\left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)' + \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)^2 - (\alpha')^2 \right) \frac{(-x \sin \alpha + y \cos \alpha)^2}{2} \\ & - \left(\left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)' + 2 \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)^2 \right) \frac{z^2}{2} \\ & + (\alpha'' + 2\alpha' \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)) (x \cos \alpha + y \sin \alpha) (-x \sin \alpha + y \cos \alpha) \\ & \left. - 12\nu \alpha' \frac{-x \sin \alpha + y \cos \alpha}{x \cos \alpha + y \sin \alpha} + \frac{12\nu^2}{(x \cos \alpha + y \sin \alpha)^2} \right), \end{aligned} \quad (1.15)$$

where α is any function of t , and a, b are arbitrary real constants. The above solution may partially reflect important features of vortices.

The paper is organized as follows. In order to get initial feeling of the MHD equations, we solve the equations in Section 2 by assuming that the solutions are linear in the spacial variables. Asymmetric approaches are presented in Section 3. In Section 4, we obtain five families solutions by the moving-frame method.

2 Linear solutions

In this section, we find four families of solutions of the MHD equations (1.1)-(1.4), which are linear with respect to the spacial variables x , y and z .

Denote

$$W_1 = u_t + uu_x + vu_y + wu_z - \mu_0(H^2(H_x^2 - H_y^1) - H^3(H_z^1 - H_x^3)) - \nu\Delta u, \quad (2.1)$$

$$W_2 = v_t + uv_x + vv_y + wv_z - \mu_0(H^3(H_y^3 - H_z^2) - H^1(H_x^2 - H_y^1)) - \nu\Delta v, \quad (2.2)$$

$$W_3 = w_t + uw_x + vw_y + ww_z - \mu_0(H^1(H_z^1 - H_x^3) - H^2(H_y^3 - H_z^2)) - \nu\Delta w. \quad (2.3)$$

To solve (1.2), we first deal with the compatibility equations

$$W_{1y} = W_{2x}, \quad W_{1z} = W_{3x}, \quad W_{2z} = W_{3y} \quad (2.4)$$

and then find p . The equations (1.4) can be written as:

$$H_t^1 - (uH^2 - vH^1)_y + (wH^1 - uH^3)_z - \eta\Delta H_1 = 0, \quad (2.5)$$

$$H_t^2 - (vH^3 - wH^2)_z + (uH^2 - vH^1)_x - \eta\Delta H_2 = 0, \quad (2.6)$$

$$H_t^3 - (wH^1 - uH^3)_x + (vH^3 - wH^2)_y - \eta\Delta H_3 = 0. \quad (2.7)$$

Let

$$\mathbf{v} = A \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad \mathbf{H} = B \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad (2.8)$$

where $A = (a_{i,j}(t))_{3 \times 3}$ and $B = (b_{i,j}(t))_{3 \times 3}$ are 3×3 matrices whose entries are functions in t . Then by (1.1) and (1.3), $\text{tr}(A) = \text{tr}(B) = 0$. Now the equations (2.4) become

$$\begin{cases} a'_{12} - a'_{21} + (a_{11} + a_{22})(a_{12} - a_{21}) + a_{13}a_{32} - a_{23}a_{31} \\ \quad + \mu_0((b_{11} + b_{22})(b_{12} - b_{21}) + b_{13}b_{32} - b_{23}b_{31}) = 0, \\ a'_{13} - a'_{31} + (a_{11} + a_{33})(a_{13} - a_{31}) + a_{23}a_{12} - a_{21}a_{32} \\ \quad + \mu_0((b_{11} + b_{33})(b_{13} - b_{31}) + b_{12}b_{23} - b_{21}b_{32}) = 0, \\ a'_{23} - a'_{32} + (a_{22} + a_{33})(a_{23} - a_{32}) + a_{13}a_{21} - a_{12}a_{31} \\ \quad + \mu_0((b_{22} + b_{33})(b_{23} - b_{32}) + b_{13}b_{21} - b_{12}b_{31}) = 0. \end{cases} \quad (2.9)$$

Moreover, the equations (1.4) are equivalent to the following system of ordinary differential equations:

$$\begin{cases} b'_{11} = a_{12}b_{21} - a_{21}b_{12} + a_{13}b_{31} - a_{31}b_{13}, \\ b'_{12} = a_{11}b_{12} - a_{12}b_{11} + a_{12}b_{22} - a_{22}b_{12} + a_{13}b_{32} - a_{32}b_{13}, \\ b'_{13} = 2a_{11}b_{13} - 2a_{13}b_{11} + a_{22}b_{13} - a_{13}b_{22} + a_{12}b_{23} - a_{23}b_{12}, \end{cases} \quad (2.10)$$

$$\begin{cases} b'_{21} = -a_{11}b_{21} + a_{21}b_{11} + a_{22}b_{21} - a_{21}b_{22} + a_{23}b_{31} - a_{31}b_{23}, \\ b'_{22} = -a_{12}b_{21} + a_{21}b_{12} + a_{23}b_{32} - a_{32}b_{23}, \\ b'_{23} = a_{11}b_{23} - a_{23}b_{11} + 2a_{22}b_{23} - 2a_{23}b_{22} - a_{13}b_{21} + a_{21}b_{13} \end{cases} \quad (2.11)$$

and

$$\begin{cases} b'_{31} = -2a_{11}b_{31} + 2a_{31}b_{11} - a_{22}b_{31} + a_{31}b_{22} - a_{21}b_{32} + a_{32}b_{21}, \\ b'_{32} = a_{11}b_{32} + a_{32}b_{11} - a_{12}b_{31} + a_{31}b_{12} - 2a_{22}b_{32} + 2a_{32}b_{22}, \\ b'_{33} = -a_{13}b_{31} + a_{31}b_{13} - a_{23}b_{32} + a_{32}b_{23}. \end{cases} \quad (2.12)$$

We consider the following special cases:

Case 1: $A = A^t$, $B = B^t$.

Easily see that the system (2.9) holds naturally. By (2.10)-(2.12), we find that B is a constant matrix and A is determined by following equation:

$$\begin{pmatrix} b_{12} & -b_{11} + b_{22} & b_{23} & -b_{32} & -b_{13} \\ 2b_{13} & b_{23} & -2b_{11} - b_{22} & b_{13} & -b_{12} \\ b_{23} & b_{13} & -b_{12} & b_{23} & -b_{11} - 2b_{22} \end{pmatrix} \begin{pmatrix} a_{11}(t) \\ a_{12}(t) \\ a_{13}(t) \\ a_{22}(t) \\ a_{23}(t) \end{pmatrix} = 0. \quad (2.13)$$

Note that above system has nontrivial solutions. They span a linear space. Since

$$p_x = -\rho W_1, \quad p_y = -\rho W_2, \quad p_z = -\rho W_3, \quad (2.14)$$

one gets

$$\begin{aligned} p = & -\frac{\rho}{2}((a'_{11} + \sum_{j=1}^3 a_{1j}^2)x^2 + (a'_{12} + \sum_{j=1}^3 a_{2j}^2)y^2 + (a'_{33} + \sum_{j=1}^3 a_{3j}^2)z^2) \\ & -\rho((a'_{12} + \sum_{j=1}^3 a_{1j}a_{2j})xy + (a'_{13} + \sum_{j=1}^3 a_{1j}a_{3j})xz + (a'_{23} + \sum_{j=1}^3 a_{2j}a_{3j})yz), \end{aligned} \quad (2.15)$$

modulo the transformation in (1.9).

Proposition 2.1. *Let B be any 3×3 constant symmetric matrix with zero trace and let A be a symmetric matrix function with zero trace determined by (2.13). Then we have the following solutions of the MHD equations (1.1)-(1.4):*

$$\mathbf{v} = A \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad \mathbf{H} = B \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad (2.16)$$

and p is given by (2.15).

Case 2: $A = A^t$, $B = -B^t$.

As in case 1, one shows that B is constant and

$$a_{11} = \frac{b_{12}^2 + b_{13}^2 - 2b_{23}^2}{3b_{12}b_{13}b_{23}}g(t), \quad a_{22} = \frac{b_{12}^2 - 2b_{13}^2 + b_{23}^2}{3b_{12}b_{13}b_{23}}g(t), \quad (2.17)$$

$$a_{12} = \frac{g(t)}{b_{12}}, \quad a_{13} = -\frac{g(t)}{b_{13}}, \quad a_{23} = \frac{g(t)}{b_{23}}, \quad (2.18)$$

where g is an arbitrary function of t .

Moreover,

$$\begin{aligned}
p = & -\frac{\rho}{2}((a'_{11} + \sum_{j=1}^3 a_{1j}^2 + 2\mu_0(b_{12}^2 + b_{13}^2))x^2 + (a'_{12} + \sum_{j=1}^3 a_{2j}^2 + 2\mu_0(b_{12}^2 + b_{23}^2))y^2 \\
& + (a'_{13} + \sum_{j=1}^3 a_{3j}^2 + 2\mu_0(b_{13}^2 + b_{23}^2))z^2) - \rho((a'_{12} + \sum_{j=1}^3 a_{1j}a_{2j} - 2\mu_0b_{13}b_{23})xy \\
& + (a'_{13} + \sum_{j=1}^3 a_{1j}a_{3j} + 2\mu_0b_{12}b_{23})xz + (a'_{23} + \sum_{j=1}^3 a_{2j}a_{3j} - 2\mu_0b_{12}b_{13})yz)
\end{aligned} \tag{2.19}$$

modulo the transformation in (1.9).

Proposition 2.2. *Let B be a 3×3 constant skew-symmetric matrix and let A be a symmetric matrix function with zero trace determined by (2.17) and (2.18). Then we have the following solutions of the MHD equations (1.1)-(1.4):*

$$\mathbf{v} = A \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad \mathbf{H} = B \begin{pmatrix} x \\ y \\ z \end{pmatrix} \tag{2.20}$$

and p is given by (2.19).

Case 3: $A = -A^t$, $B = B^t$.

One shows that A is constant and

$$(b_{11}, b_{12}, b_{13}, b_{22}, b_{23})^t = \exp(Mt)\mathbf{c} \tag{2.21}$$

by (2.9)-(2.12), where $\mathbf{c}$ is any five-dimensional constant vector and

$$M = \begin{pmatrix} 0 & 2a_{12} & 2a_{13} & 0 & 0 \\ -2a_{12} & 0 & a_{23} & 2a_{12} & a_{13} \\ -2a_{13} & -a_{23} & 0 & -a_{13} & a_{12} \\ 0 & -2a_{12} & 0 & 0 & 2a_{23} \\ -a_{23} & -a_{13} & -a_{12} & -2a_{23} & 0 \end{pmatrix}. \tag{2.22}$$

Moreover,

$$\begin{aligned}
p = & -\rho(-(a_{12}^2 + a_{13}^2)\frac{x^2}{2} - (a_{12}^2 + a_{23}^2)\frac{y^2}{2} - (a_{13}^2 + a_{23}^2)\frac{z^2}{2} \\
& - a_{13}a_{32}xy - a_{12}a_{23}xz - a_{21}a_{13}yz)
\end{aligned} \tag{2.23}$$

modulo the transformation in (1.9).

Proposition 2.3. *Let A be a 3×3 constant skew-symmetric matrix and let B be a symmetric matrix function with zero trace determined by (2.21). Then we have the following solutions of the MHD equations (1.1)-(1.4):*

$$\mathbf{v} = A \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad \mathbf{H} = B \begin{pmatrix} x \\ y \\ z \end{pmatrix} \tag{2.24}$$

and p is given by (2.23).

Case 4: $A = -A^t$, $B = -B^t$.

Similarly, we set

$$N = \begin{pmatrix} 0 & a_{23} & -a_{13} \\ -a_{23} & 0 & a_{12} \\ a_{13} & -a_{12} & 0 \end{pmatrix} \quad (2.25)$$

and

$$(b_{12}, b_{13}, b_{23})^t = \exp(Nt)\mathbf{c}, \quad (2.26)$$

where $\mathbf{c}$ is a three-dimensional constant vector. Then we get the following proposition:

Proposition 2.4. *Let A be a 3×3 constant skew-symmetric matrix and let B be a skew-symmetric matrix function determined by (2.26). Then we have the following solutions of the MHD equations (1.1)-(1.4):*

$$\mathbf{v} = A \begin{pmatrix} x \\ y \\ z \end{pmatrix}, \quad \mathbf{H} = B \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad (2.27)$$

and

$$\begin{aligned} p = & -\rho((-a_{12}^2 + a_{13}^2) + 2\mu_0(b_{12}^2 + b_{13}^2))\frac{x^2}{2} + (-a_{12}^2 + a_{23}^2) + 2\mu_0(b_{12}^2 + b_{23}^2))\frac{y^2}{2} \\ & + (-a_{13}^2 + a_{23}^2) + 2\mu_0(b_{13}^2 + b_{23}^2))\frac{z^2}{2} + (-a_{13}a_{32} + 2\mu_0b_{13}b_{32})xy \\ & + (-a_{12}a_{23} + 2\mu_0b_{12}b_{23})xz + (-a_{21}a_{13} + 2\mu_0b_{21}b_{13})yz. \end{aligned} \quad (2.28)$$

3 Asymmetric Approaches

In this section, we will solve the MHD equations (1.1)-(1.4) by imposing asymmetric assumptions on $\mathbf{v}$ and $\mathbf{H}$, following [32].

Firstly, we set

$$\begin{cases} u = f + (\beta - \psi_z)x, \\ v = -\beta y, \\ w = \psi \end{cases} \quad \text{and} \quad \begin{cases} H^1 = g + (\gamma - \varphi_z)x, \\ H^2 = -\gamma y, \\ H^3 = \varphi, \end{cases} \quad (3.1)$$

where f and g are functions of t , y and z ; φ and ψ are functions of t and z ; β and γ are functions of t .

By (2.1)-(2.3),

$$\begin{aligned} W_1 = & f_t + f(\beta - \psi_z) - \beta y f_y + \psi f_z - \nu(f_{yy} + f_{zz}) - \mu_0(\gamma y g_y - \varphi g_z) \\ & + ((\beta - \psi_z)^2 + \beta' - \psi_{zt} - \psi \psi_{zz} + \nu \psi_{zzz} - \mu_0 \varphi \varphi_{zz})x, \end{aligned} \quad (3.2)$$

$$W_2 = (\beta^2 + \beta')y - \mu_0 g_y g - \mu_0 g_y (\gamma - \varphi_z)x, \quad (3.3)$$

and

$$W_3 = \psi_z(1 + \psi) - \nu \psi_{zz} - \mu_0(g_z g + (g_z(\gamma - \varphi_z) - g \varphi_{zz})x - \varphi_{zz}(\gamma - \varphi_z)x^2). \quad (3.4)$$

According to (1.4) and (2.4), we find that $\gamma = b/2$ is a constant and

$$\begin{cases} \varphi_t = -\psi \varphi_z + \psi_z \varphi + \eta \varphi_{zz}, \\ g_t = -\frac{b}{2}(y f_y - f) + \beta(g + y g_y) - \psi_z g - \psi g_z + f \varphi_z + f_z \varphi + \eta(g_{yy} + g_{zz}), \end{cases} \quad (3.5)$$

$$\begin{cases} f_{yt} - \psi_z f_y - \beta y f_{yy} + \psi f_{yz} - \nu(f_{yyy} + f_{zzy}) - \mu_0(\frac{b}{2}y g_{yy} + g_y \varphi_z - \varphi g_{yz}) = 0, \\ f_{zt} + f_z \beta - f \psi_{zz} - \beta y f_{yz} + \psi f_{zz} - \nu(f_{yzz} + f_{zzz}) - \mu_0(\frac{b}{2}y g_{yz} - \varphi g_{zz} - \frac{b}{2}g_z + g \varphi_{zz}) = 0, \end{cases} \quad (3.6)$$

$$-2\beta\psi_{zz} + \psi_z\psi_{zz} - \psi_{zzt} - \psi\psi_{zzz} + \nu\psi_{zzzz} - \mu_0(b - \varphi_z\varphi_{zz} + \varphi\varphi_{zzz}) = 0. \quad (3.7)$$

One shows that the system (3.6) is implied by

$$f_t + (\beta - \psi_z)f - \beta y f_y + \psi f_z - \nu(f_{yy} + f_{zz}) - \mu_0\left(\frac{b}{2}yg_y - \varphi g_z + g\left(\frac{b}{2} - \varphi_z\right)\right) = 0. \quad (3.8)$$

We assume that

$$\varphi = az + l. \quad (3.9)$$

Then

$$\psi = ahz + hl - \frac{l'}{a} \quad (3.10)$$

by (3.5)₁, where a is a constant and h, l are functions of t . Hence the equations (3.5)-(3.7) can be written as

$$\begin{cases} f_t + (\beta - ah)f - \beta y f_y + (h(az + l) - \frac{l'}{a})f_z - \mu_0\left(\frac{b}{2}yg_y - (az + l)g_z - \left(\frac{b}{2} - a\right)g\right) \\ - \nu(f_{yy} + f_{zz}) = 0, \\ g_t + \frac{b}{2}(yf_y + f) - (\beta - ah)g - \beta yg_y + (ahz + hl - \frac{l'}{a})g_z - af - (az + l)f_z - \eta(g_{yy} + g_{zz}) = 0. \end{cases} \quad (3.11)$$

Now we solve above equations. Firstly we suppose that l is a constant and then use the method of separation of variables. Set

$$f = y(az + l)s, \quad g = y(az + l)q, \quad (3.12)$$

where s and q are functions of t . Substituting (3.12) into (3.11), one gets that s is a constant and

$$q = \exp\left(2 \int (\beta - ah)dt\right) \left((2a - b)s \int \exp\left(-2 \int (\beta - ah)dt\right)dt + c\right), \quad (3.13)$$

where c is a constant. Furthermore,

$$\begin{aligned} p = & -\rho((\beta' - ah' + (\beta - ah)^2)\frac{x^2}{2} + (-\beta' + \beta^2)\frac{y^2}{2} + \frac{1}{2a}(h' + ah^2)a^2z^2 \\ & - \mu_0q^2a^2z^2\frac{y^2}{2} - \mu_0qa\left(\frac{b}{2} - a\right)xyz) \end{aligned} \quad (3.14)$$

modulo the transformations in (1.8) and in (1.9).

Theorem 3.1. *For any functions $h(t)$, $\beta(t)$ and constants a, b, c , and s , we have the following solutions of the MHD equations (1.1)-(1.4):*

$$\begin{cases} u = asyz + (\beta - ah)x, \\ v = -\beta y, \\ w = ahz, \end{cases} \quad (3.15)$$

$$\begin{cases} H^1 = ayz \exp\left(2 \int (\beta - ah)dt\right) \left((2a - b)s \int \exp\left(-2 \int (\beta - ah)dt\right)dt + c\right) + \left(\frac{b}{2} - a\right)x, \\ H^2 = -\frac{b}{2}y, \\ H^3 = az \end{cases} \quad (3.16)$$

and p is given by (3.14).

Now we give another solutions of (3.11). Motivated from (3.12), we assume that

$$\begin{cases} f = Cy^2 + Dz^2 + Eyz + Fy + Gz + H, \\ g = Iy^2 + Jz^2 + Kyz + Ly + Mz + N, \end{cases} \quad (3.17)$$

where $C, D, E, F, G, H, I, J, K, L, M$ and N are functions of t .

Substituting (3.17) into (3.11), we get that

$$\begin{cases} C' - (\beta + ah)C - \mu_0(\frac{b}{2} + a)I = 0, \\ D' + (\beta + ah)D + \mu_0(\frac{b}{2} + a)J = 0, \\ E' = 0, \\ F' - ahF + (hl - \frac{l'}{a})E - a\mu_0L + \mu_0lK = 0, \\ G' + \beta G + 2(hl - \frac{l'}{a})D + \frac{b}{2}\mu_0M + 2\mu_0lJ = 0, \\ H' + (\beta - ah)H + (hl - \frac{l'}{a})G - 2\nu(C + E) + \mu_0lM + \mu_0(\frac{b}{2} - a)N = 0 \end{cases} \quad (3.18)$$

and

$$\begin{cases} I' - (3\beta - ah)I + (\frac{3b}{2} - a)C = 0, \\ J' - (\beta - 3ah)J + (\frac{b}{2} - 3a)D = 0, \\ K' - 2(\beta - ah)K + 2(b - 2a)E = 0, \\ L' - (2\beta - ah)L + (b - a)F + (hl - \frac{l'}{a})K - lE = 0, \\ M' - (\beta - 2ah)M + (\frac{b}{2} - 2a)G + 2(hl - \frac{l'}{a})J - 2lD = 0, \\ N' - (\beta - ah)N + (\frac{b}{2} - a)H + (hl - \frac{l'}{a})M - 2\eta(I + J) - lG = 0. \end{cases} \quad (3.19)$$

Thus we have that

$$\begin{cases} I = \frac{1}{(\frac{b}{2} + a)\mu_0}(C' - (\beta + ah)C), \\ C'' - 4\beta C' - (\beta' + ah' - (3\beta - ah)(\beta + ah) - (\frac{3b}{2} - a)(\frac{b}{2} + a)\mu_0)C = 0, \end{cases} \quad (3.20)$$

$$\begin{cases} J = -\frac{1}{(\frac{b}{2} + a)\mu_0}(D' + (\beta + ah)D), \\ D'' + 4ahD' + (\beta' + ah' - (\beta - 3ah)(\beta + ah) - (\frac{b}{2} - 3a)(\frac{b}{2} + a)\mu_0)D = 0, \end{cases} \quad (3.21)$$

$$\begin{cases} E' = 0, \\ K' - 2(\beta - ah)K = -(b - 2a)E, \end{cases} \quad (3.22)$$

$$\begin{cases} L = \frac{1}{a\mu_0}(F' - ahF + (hl - \frac{l'}{a})E + \mu_0lK), \\ F'' - 2\beta F' + (ah(2\beta - ah) - ah' + (b - a)a\mu_0)F = f_F(t), \end{cases} \quad (3.23)$$

$$\begin{cases} M = -\frac{2}{b\mu_0}(G' + \beta G + 2(hl - \frac{l'}{a})D + 2\mu_0lJ), \\ G'' + 2ahG' + (\beta' - \beta(\beta - 2ah) - \frac{b}{2}(\frac{b}{2} - 2a)\mu_0)G = f_G(t) \end{cases} \quad (3.24)$$

and

$$\begin{cases} N = -\frac{1}{(\frac{b}{2} - a)\mu_0}(H' + (\beta - ah)H + (hl - \frac{l'}{a})G - 2\nu(C + E) + \mu_0lM), \\ H'' + (\beta' - ah' - (\beta - ah)^2 - (\frac{b}{2} - a)^2\mu_0)H = f_H(t), \end{cases} \quad (3.25)$$

where

$$f_F(t) = -((hl - \frac{l'}{a})E + \mu_0 lK)' + 2(\beta - ah)((hl - \frac{l'}{a})E + \mu_0 lK) - a\mu_0((hl - \frac{l'}{a})K - lE), \quad (3.26)$$

$$f_G(t) = -2((hl - \frac{l'}{a})D + \mu_0 lJ)' + 2(\beta - 2ah)((hl - \frac{l'}{a})D + \mu_0 lJ) + b\mu_0((hl - \frac{l'}{a})J - lD), \quad (3.27)$$

and

$$\begin{aligned} f_H(t) = & -(hl - \frac{l'}{a})G + 2\nu(C + E) - \mu_0 lM)' + (\beta - ah)((hl - \frac{l'}{a})G - 2\nu(C + E) + \mu_0 lM) \\ & + (\frac{b}{2} - a)\mu_0((hl - \frac{l'}{a})M - 2\eta(I + J) - lG). \end{aligned} \quad (3.28)$$

To write down the solutions of above systems explicitly, we assume

$$\beta - ah = q, \quad (3.29)$$

where q is a constant. Under this condition, the equations $(3.20)_2$, $(3.21)_2$, $(3.23)_2$ and $(3.24)_2$ can be written as the following form via certain changing of variables:

$$A''(t) + kA(t) = f_A(t). \quad (3.30)$$

Here k is a constant. We will explain this by solving $(3.20)_2$ in detail. Since the other equations are similar, we will only write down the solutions but omit the solving procedure.

Set

$$C = \exp(2 \int \beta dt) \tilde{C}. \quad (3.31)$$

Then $(3.20)_2$ becomes

$$\tilde{C}'' - (q^2 - (\frac{3b}{2} - a)(\frac{b}{2} + a)\mu_0)\tilde{C} = 0. \quad (3.32)$$

We write $\lambda_C = q^2 - (\frac{3b}{2} - a)(\frac{b}{2} + a)\mu_0$ and define

$$\xi_{C_1} = \begin{cases} \exp(\sqrt{\lambda_C} t + 2 \int \beta dt) & \text{if } \lambda_C > 0, \\ \exp(2 \int \beta dt) & \text{if } \lambda_C = 0, \\ \cos(i\sqrt{\lambda_C} t) \exp(2 \int \beta dt) & \text{if } \lambda_C < 0, \end{cases} \quad (3.33)$$

$$\xi_{C_2} = \begin{cases} \exp(-\sqrt{\lambda_C} t + 2 \int \beta dt) & \text{if } \lambda_C > 0, \\ t \exp(2 \int \beta dt) & \text{if } \lambda_C = 0, \\ \sin(i\sqrt{\lambda_C} t) \exp(2 \int \beta dt) & \text{if } \lambda_C < 0. \end{cases} \quad (3.34)$$

Then

$$C = l_{C_1} \xi_{C_1} + l_{C_2} \xi_{C_2} \quad (3.35)$$

and

$$I = \frac{\sum_{j=1}^2 l_{C_j} (\xi'_{C_j} - (2\beta - q)\xi_{C_j})}{(\frac{b}{2} + a)\mu_0}, \quad (3.36)$$

where l_{C_1} and l_{C_2} are real constants.

Similarly, we set $\lambda_D = q^2 + (\frac{b}{2} - 3a)(\frac{b}{2} + a)\mu_0$ and define

$$\xi_{D_1} = \begin{cases} \exp(\sqrt{\lambda_D} t - 2 \int (\beta + q) dt) & \text{if } \lambda_D > 0, \\ \exp(-2 \int (\beta + q) dt) & \text{if } \lambda_D = 0, \\ \cos(i\sqrt{\lambda_D} t) \exp(-2 \int (\beta + q) dt) & \text{if } \lambda_D < 0, \end{cases} \quad (3.37)$$

$$\xi_{D_2} = \begin{cases} \exp(-\sqrt{\lambda_D} t - 2 \int (\beta + q) dt) & \text{if } \lambda_D > 0, \\ t \exp(-2 \int (\beta + q) dt) & \text{if } \lambda_D = 0, \\ \sin(i\sqrt{\lambda_D} t) \exp(-2 \int (\beta + q) dt) & \text{if } \lambda_D < 0. \end{cases} \quad (3.38)$$

Then

$$D = l_{D_1}\xi_{D_1} + l_{D_2}\xi_{D_2} \quad (3.39)$$

and

$$J = -\frac{\sum_{j=1}^2 l_{D_j}(\xi'_{D_j} + (2\beta - q)\xi_{D_j})}{(\frac{b}{2} + a)\mu_0}, \quad (3.40)$$

where l_{D_1} and l_{D_2} are real constants.

Given functions $r(t)$ and $s(t)$, we denote their Wronskian determinant by

$$W[r(t), s(t)] = \det \begin{pmatrix} r(t) & s(t) \\ r'(t) & s'(t) \end{pmatrix} = r(t)s'(t) - r'(t)s(t). \quad (3.41)$$

For notational convenience, we write

$$W_1[r(t), s(t)] = \det \begin{pmatrix} 0 & s(t) \\ 1 & s'(t) \end{pmatrix} = -s(t), \quad (3.42)$$

$$W_2[r(t), s(t)] = \det \begin{pmatrix} r(t) & 0 \\ r'(t) & 1 \end{pmatrix} = r(t). \quad (3.43)$$

Set $\lambda_F = q^2 - (b - a)a\mu_0$ and define

$$\xi_{F_1} = \begin{cases} \exp(\sqrt{\lambda_F} t + \int \beta dt) & \text{if } \lambda_F > 0, \\ \exp(\int \beta dt) & \text{if } \lambda_F = 0, \\ \cos(i\sqrt{\lambda_F} t) \exp(\int \beta dt) & \text{if } \lambda_F < 0, \end{cases} \quad (3.44)$$

$$\xi_{F_2} = \begin{cases} \exp(-\sqrt{\lambda_F} t + \int \beta dt) & \text{if } \lambda_F > 0, \\ t \exp(\int \beta dt) & \text{if } \lambda_F = 0, \\ \sin(i\sqrt{\lambda_F} t) \exp(\int \beta dt) & \text{if } \lambda_F < 0. \end{cases} \quad (3.45)$$

Then

$$F = \sum_{j=1}^2 \xi_{F_j}(l_{F_j} + \int_0^t \frac{W_j[\xi_{F_1}(s), \xi_{F_2}(s)]}{W[\xi_{F_1}(s), \xi_{F_2}(s)]} f_F(s) ds) \quad (3.46)$$

and

$$\begin{aligned} L = & \frac{1}{a\mu_0} \left(\frac{1}{a} ((\beta - q)l - l' + \frac{a(b - 2a)}{2q})E + ke^{2qt} + \sum_{j=1}^2 (\xi_{F_j} \frac{W_j[\xi_{F_1}(t), \xi_{F_2}(t)]}{W[\xi_{F_1}(t), \xi_{F_2}(t)]} f_F(t) \right. \\ & \left. + (\xi'_{F_j} - (\beta - q)\xi_{F_j})(l_{F_j} + \int_0^t \frac{W_j[\xi_{F_1}(s), \xi_{F_2}(s)]}{W[\xi_{F_1}(s), \xi_{F_2}(s)]} f_F(s) ds) \right), \end{aligned} \quad (3.47)$$

where l_{F_1} and l_{F_2} are real constants. In this expression, we have use the fact:

$$K = ke^{2qt} + \frac{b - 2a}{2q}E, \quad (3.48)$$

where k and E are real constant.

Let $\lambda_G = q^2 + \frac{b}{2}(\frac{b}{2} - 2a)\mu_0$ and denote

$$\xi_{G_1} = \begin{cases} \exp(\sqrt{\lambda_G} t - \int (\beta - q) dt) & \text{if } \lambda_G > 0, \\ \exp(-\int (\beta - q) dt) & \text{if } \lambda_G = 0, \\ \cos(i\sqrt{\lambda_G} t) \exp(-\int (\beta - q) dt) & \text{if } \lambda_G < 0, \end{cases} \quad (3.49)$$

$$\xi_{G_2} = \begin{cases} \exp(-\sqrt{\lambda_G} t - \int (\beta - q) dt) & \text{if } \lambda_G > 0, \\ t \exp(-\int (\beta - q) dt) & \text{if } \lambda_G = 0, \\ \sin(i\sqrt{\lambda_G} t) \exp(-\int (\beta - q) dt) & \text{if } \lambda_G < 0. \end{cases} \quad (3.50)$$

Then

$$G = \sum_{j=1}^2 \xi_{G_j} (l_{G_j} + \int_0^t \frac{W_j[\xi_{G_1}(s), \xi_{G_2}(s)]}{W[\xi_{G_1}(s), \xi_{G_2}(s)]} f_G(s) ds) \quad (3.51)$$

and

$$\begin{aligned} M = & -\frac{2}{\mu_0} \left(\sum_{j=1}^2 (2l_{D_j} \left(\frac{(\beta - q - l') \xi_{D_j}}{a} - \frac{l}{\frac{b}{2} + a} (\xi'_{D_j} + (2\beta - q) \xi_{D_j}) \right) \right. \\ & + (\xi'_{G_j} + \beta \xi_{G_j}) (l_{G_j} + \int_0^t \frac{W_j[\xi_{G_1}(s), \xi_{G_2}(s)]}{W[\xi_{G_1}(s), \xi_{G_2}(s)]} f_G(s) ds) \\ & \left. + \xi_{G_j} \frac{W_j[\xi_{G_1}(t), \xi_{G_2}(t)]}{W[\xi_{G_1}(t), \xi_{G_2}(t)]} f_G(t) \right), \end{aligned} \quad (3.52)$$

where l_{G_1} and l_{G_2} are real constants.

Set $\lambda_H = q^2 + (\frac{b}{2} - a)\mu_0^2$ and define

$$\xi_{H_1} = \begin{cases} \exp(\sqrt{\lambda_H} t) & \text{if } \lambda_H > 0, \\ 1 & \text{if } \lambda_H = 0, \\ \cos(i\sqrt{\lambda_H} t) & \text{if } \lambda_H < 0, \end{cases} \quad (3.53)$$

$$\xi_{H_2} = \begin{cases} \exp(-\sqrt{\lambda_H} t) & \text{if } \lambda_H > 0, \\ t & \text{if } \lambda_H = 0, \\ \sin(i\sqrt{\lambda_H} t) & \text{if } \lambda_H < 0. \end{cases} \quad (3.54)$$

Then

$$H = \sum_{j=1}^2 \xi_{H_j} (l_{H_j} + \int_0^t \frac{W_j[\xi_{H_1}(s), \xi_{H_2}(s)]}{W[\xi_{H_1}(s), \xi_{H_2}(s)]} f_H(s) ds) \quad (3.55)$$

and

$$\begin{aligned} N = & -\frac{1}{(\frac{b}{2} - a)\mu_0} \left(\frac{1}{a} ((\beta - q)l - l')G - 2\nu(C + E) + \mu_0 lM + \sum_{j=1}^2 (\xi_{H_j} \frac{W_j[\xi_{H_1}(t), \xi_{H_2}(t)]}{W[\xi_{H_1}(t), \xi_{H_2}(t)]} f_H(t) \right. \\ & \left. + (\xi'_{H_j} + q\xi_{H_j}) (l_{H_j} + \int_0^t \frac{W_j[\xi_{H_1}(s), \xi_{H_2}(s)]}{W[\xi_{H_1}(s), \xi_{H_2}(s)]} f_H(s) ds) \right), \end{aligned} \quad (3.56)$$

where E , l_{H_1} and l_{H_2} are real constants, C , G and M are given by (3.35), (3.51) and (3.52).

Furthermore, by (2.14), one has

$$\begin{aligned} p = & -\rho \left(\frac{q^2}{2} x^2 + (H' + qH + \frac{(\beta - q)l - l'}{a} G + \mu_0 lM - 2\nu(C + D))x \right. \\ & - \frac{\mu_0}{2} I^2 y^4 - \mu_0 I L y^3 + \frac{1}{2} (-\beta' + \beta^2 - \mu_0(2IN + L^2)) y^2 - \mu_0 L N y \\ & - \frac{\mu_0}{2} J^2 z^4 - \mu_0 J M z^3 + (\beta' + (\beta - q)^2 - \mu_0(2JN + M^2)) z^2 \\ & + (-\mu_0 M N + \frac{1}{a} (\beta' l + (\beta - q)l' - l'') + \frac{\beta - q}{a} ((\beta - q)l - l')) z \\ & - (\frac{b}{2} - a)\mu_0 K x y z - \frac{b - 2a}{2} \mu_0 I x y^2 - (\frac{b}{2} - a)\mu_0 L x y - (\frac{b}{2} - a)\mu_0 J x z^2 - (\frac{b}{2} - a)\mu_0 M x z \\ & - \mu_0 K J y z^3 - \mu_0 K I y^3 z - \frac{\mu_0}{2} (2IJ + K^2) y^2 z^2 - \mu_0 (JL + KM) y z^2 \\ & \left. - \mu_0 (KL + IM) y^z - \mu_0 (KN + LM) y z \right) \end{aligned} \quad (3.57)$$

modulo the transformation in (1.9).

Theorem 3.2. For any functions $\beta(t)$, $l(t)$ and real constants a , b , q , E , we have the following solutions of the MHD equations (1.1)-(1.4):

$$\begin{cases} u = Cy^2 + Dz^2 + Eyz + Fy + Gz + H + qx, \\ v = -\beta y, \\ w = (\beta - q)z + \frac{(\beta - q)l - l'}{a}, \end{cases} \quad (3.58)$$

$$\begin{cases} H^1 = Iy^2 + Jz^2 + Kyz + Ly + Mz + N + \left(\frac{b}{2} - a\right)x, \\ H^2 = -\frac{b}{2}y, \\ H^3 = az + l \end{cases} \quad (3.59)$$

and p is given by (3.57). Here the functions C , D , F , G , H , I , J , K , L , M and N are given by (3.35), (3.39), (3.46), (3.51), (3.55), (3.36), (3.40), (3.48), (3.47), (3.52) and (3.56), respectively.

We denote

$$\varpi = x^2 + y^2 \quad (3.60)$$

and assume

$$\begin{cases} u = \frac{\alpha}{2}x + y\phi(t, \varpi), \\ v = \frac{\alpha}{2}y - x\phi(t, \varpi), \\ w = -\alpha z + \psi(t, \varpi), \end{cases} \quad \begin{cases} H^1 = \frac{\beta}{2}x + y\xi(t, \varpi), \\ H^2 = \frac{\beta}{2}y - x\xi(t, \varpi), \\ H^3 = -\beta z + \sigma(t, \varpi). \end{cases} \quad (3.61)$$

Recall that (1.4) is just

$$\begin{cases} H_t^1 = (uH^2 - vH^1)_y - (wH^1 - uH^3)_z + \eta\Delta H^1, \\ H_t^2 = (vH^3 - wH^2)_z - (uH^2 - vH^1)_x + \eta\Delta H^2, \\ H_t^3 = (wH^1 - uH^3)_x - (vH^3 - wH^2)_y + \eta\Delta H^3. \end{cases} \quad (3.62)$$

Substituting (3.61) into (3.62)₁, one gets that $\beta = b$ is a constant and

$$\xi_t = -\alpha\varpi\xi_\varpi + b\varpi\phi_\varpi + 4\eta(\varpi\xi)_{\varpi\varpi}. \quad (3.63)$$

One shows that (3.62)₂ is equivalent to (3.63), and (3.62)₃ implies that

$$\sigma_t = b(\varpi\psi)_\varpi - \alpha\varpi\sigma_\varpi + 4\eta(\sigma_\varpi + \varpi\sigma_{\varpi\varpi}). \quad (3.64)$$

Moreover, by (2.4), we have

$$\begin{cases} (\varpi\phi)_{t\varpi} + \alpha(\varpi(\varpi\phi)_\varpi)_\varpi - 4\nu(\varpi(\varpi\phi)_{\varpi\varpi})_\varpi + b\mu_0(\varpi(\varpi\xi)_\varpi)_\varpi = 0, \\ \psi_t - \alpha\psi + \alpha\varpi\psi_\varpi - 4\nu(\psi_\varpi + \varpi\psi_{\varpi\varpi}) + b\mu_0\varpi\sigma_\varpi - b\mu_0\sigma = 0. \end{cases} \quad (3.65)$$

Given any $n \in \mathbb{N}$, set

$$\phi = \sum_{m=0}^n a_m(t)\varpi^m, \quad \xi = \sum_{m=0}^n c_m(t)\varpi^m, \quad \sigma = \sum_{m=0}^n f_m(t)\varpi^m, \quad \psi = \sum_{m=0}^n g_m(t)\varpi^m. \quad (3.66)$$

Substituting (3.66) into (3.63)-(3.65) and considering the coefficients of ϖ^m , we obtain that

$$\begin{cases} a_{m,t} + (m+1)\alpha a_m + b\mu_0(m+1)c_m = 4\nu(m+1)(m+2)a_{m+1}, \\ c_{m,t} - bma_m + \alpha mc_m = 4\eta(m+1)(m+2)c_{m+1} \end{cases} \quad (3.67)$$

and

$$\begin{cases} f_{m,t} = -\alpha m f_m + b(m+1)g_m + 4\eta(m+1)^2 f_{m+1}, \\ g_{m,t} = -(m-1)\alpha g_m - b(m-1)\mu_0 f_m + 4\nu(m+1)^2 g_{m+1}, \end{cases} \quad (3.68)$$

where $a_{-k} = a_{n+k} = c_{-k} = c_{n+k} = f_{-k} = f_{n+k} = g_{-k} = g_{n+k} = 0$ for $k > 0$. These systems are similar to (3.20). If we set

$$\alpha = \frac{k + 2k^2 l e^{-kt}}{-1 + 2kl e^{-kt}} \quad (3.69)$$

for any constants k and l , then we can transform them into certain second order ordinary differential equations with constant coefficients, as we did in (3.20).

Observe that

$$c_m = -\frac{1}{(m+1)b\mu_0}(a_{m,t} + (m+1)\alpha a_m - 4\nu(m+1)(m+2)a_{m+1}). \quad (3.70)$$

Substituting (3.70) into (3.67), one has

$$a_m'' + (2m+1)\alpha a_m' + (m+1)(m\alpha^2 + \alpha' + b^2\mu_0 m)a_m = F_m(t), \quad (3.71)$$

where

$$F_m = 4(m+1)(m+2)(m\nu\alpha a_{m+1} + \nu a_{m+1}' + \eta c_{m+1}). \quad (3.72)$$

Set

$$a_m = \tilde{a}_m \exp\left(-\frac{1}{2} \int (2m+1)\alpha dt\right) = (e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt})\tilde{a}_m. \quad (3.73)$$

Then (3.71) becomes

$$\tilde{a}_m'' + ((m+1)(m\alpha^2 + \alpha' + b^2\mu_0 m) - \frac{2m+1}{2}\alpha' - \frac{1}{4}(2m+1)^2\alpha^2)\tilde{a}_m = F_m(t). \quad (3.74)$$

By (3.69),

$$(m+1)(m\alpha^2 + \alpha' + b^2\mu_0 m) - \frac{2m+1}{2}\alpha' - \frac{1}{4}(2m+1)^2\alpha^2 = m(m+1)b^2\mu_0 - \frac{1}{4}k^2 \quad (3.75)$$

is a constant. Let $\lambda_m = -m(m+1)b^2\mu_0 + \frac{1}{4}k^2$ and define

$$\zeta_{m,1} = \begin{cases} \exp(\sqrt{\lambda_m} t)(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \lambda_m > 0, \\ e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt} & \text{if } \lambda_m = 0, \\ \cos(i\sqrt{\lambda_m} t)(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \lambda_m < 0, \end{cases} \quad (3.76)$$

$$\zeta_{m,2} = \begin{cases} \exp(-\sqrt{\lambda_m} t)(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \lambda_m > 0, \\ t(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \lambda_m = 0, \\ \sin(i\sqrt{\lambda_m} t)(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \lambda_m < 0. \end{cases} \quad (3.77)$$

Then we can get a_m and c_m by the following recursive formulae

$$a_m = \sum_{j=1}^2 \zeta_{m,j}(l_{m,j} + \int_0^t \frac{W_j[\zeta_{m,1}(s), \zeta_{m,2}(s)]}{W[\zeta_{m,1}(s), \zeta_{m,2}(s)]} F_m(s) ds), \quad (3.78)$$

$$\begin{aligned} c_m = & -\frac{1}{b\mu_0(m+1)} \sum_{j=1}^2 \left(\zeta_{m,j}'(l_{m,j} + \int_0^t \frac{W_j[\zeta_{m,1}(s), \zeta_{m,2}(s)]}{W[\zeta_{m,1}(s), \zeta_{m,2}(s)]} F_m(s) ds) \right. \\ & + \zeta_{m,j} \left(\frac{W_j[\zeta_{m,1}(t), \zeta_{m,2}(t)]}{W[\zeta_{m,1}(t), \zeta_{m,2}(t)]} F_m(t) + (m+1)\alpha(l_{m,j} + \int_0^t \frac{W_j[\zeta_{m,1}(s), \zeta_{m,2}(s)]}{W[\zeta_{m,1}(s), \zeta_{m,2}(s)]} F_m(s) ds) \right. \\ & \left. \left. + 4\nu(m+1)(m+2)\zeta_{m+1,j}(l_{m+1,j} + \int_0^t \frac{W_j[\zeta_{m+1,1}(s), \zeta_{m+1,2}(s)]}{W[\zeta_{m+1,1}(s), \zeta_{m+1,2}(s)]} F_{m+1}(s) ds) \right) \right), \quad (3.79) \end{aligned}$$

where $l_{m,1}$ and $l_{m,2}$ are real constants.

Given

$$G_m(t) = 4(m+1)^2((m-1)\alpha\eta f_{m+1} + \eta f'_{m+1} + \nu g_{m+1}), \quad (3.80)$$

the solutions of system (3.68) can be written down similarly as those of (3.78) and (3.79).

Set

$$\tau_m = \frac{1}{4}k^2 - (m^2 - 1)b^2\mu_0 \quad (3.81)$$

and denote

$$\varsigma_{m,1} = \begin{cases} \exp(\sqrt{\tau_m} t)(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \tau_m > 0, \\ e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt} & \text{if } \tau_m = 0, \\ \cos(i\sqrt{\tau_m} t)(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \tau_m < 0, \end{cases} \quad (3.82)$$

$$\varsigma_{m,2} = \begin{cases} \exp(-\sqrt{\tau_m} t)(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \tau_m > 0, \\ t(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \tau_m = 0, \\ \sin(i\sqrt{\tau_m} t)(e^{(m+\frac{1}{2})kt} - 2kle^{(m-\frac{1}{2})kt}) & \text{if } \tau_m < 0. \end{cases} \quad (3.83)$$

Then

$$f_m = \sum_{j=1}^2 \varsigma_{m,j}(q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} G_m(s) ds) \quad (3.84)$$

and

$$\begin{aligned} g_m = & \frac{1}{b(m+1)} \sum_{j=1}^2 \left(\varsigma'_{m,j}(q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} G_m(s) ds) \right. \\ & + \varsigma_{m,j} \left(\frac{W_j[\varsigma_{m,1}(t), \varsigma_{m,2}(t)]}{W[\varsigma_{m,1}(t), \varsigma_{m,2}(t)]} G_m(t) + m\alpha(q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} G_m(s) ds) \right. \\ & \left. \left. - 4\eta(m+1)^2 \varsigma_{m+1,j}(q_{m+1,j} + \int_0^t \frac{W_j[\varsigma_{m+1,1}(s), \varsigma_{m+1,2}(s)]}{W[\varsigma_{m+1,1}(s), \varsigma_{m+1,2}(s)]} G_{m+1}(s) ds) \right) \right), \end{aligned} \quad (3.85)$$

where $q_{m,1}$ and $q_{m,2}$ are real constants. The function p can be written as

$$p = -\rho \left(\frac{\alpha' + \alpha^2}{2} \varpi + \frac{1}{2} (2\alpha^2 - \alpha') z^2 + b\mu_0 z\sigma - \frac{1}{2} \int \phi^2 d\varpi - \xi^2 + \int \varpi \xi \xi_\varpi d\varpi \right) \quad (3.86)$$

modulo the transformation in (1.9).

Theorem 3.3. *Given any constant b , positive integer n and the function*

$$\alpha = \frac{k + 2k^2 l e^{-kt}}{-1 + 2k l e^{-kt}} \quad (3.87)$$

with constants k and l , we have the following solutions of the MHD equations (1.1)-(1.4):

$$\begin{cases} u = \frac{\alpha}{2}x + y \sum_{m=0}^n a_m(t)(x^2 + y^2)^m, \\ v = \frac{\alpha}{2}y - x \sum_{m=0}^n a_m(t)(x^2 + y^2)^m, \\ w = -\alpha z + \sum_{m=0}^n g_m(t)(x^2 + y^2)^m, \end{cases} \quad \begin{cases} H^1 = \frac{b}{2}x + y \sum_{m=0}^n c_m(t)(x^2 + y^2)^m, \\ H^2 = \frac{b}{2}y - x \sum_{m=0}^n c_m(t)(x^2 + y^2)^m, \\ H^3 = -bz + \sum_{m=0}^n f_m(t)(x^2 + y^2)^m. \end{cases} \quad (3.88)$$

and p is given by (3.86). Here the functions a_m , c_m , f_m and g_m are given by (3.78), (3.79), (3.84) and (3.85), respectively.

4 Moving-Frame Approach

In this section, we use the moving-frame method to find six families exact solutions of the MHD equations (1.1)-(1.4).

Let α and β be two functions in t . We denote

$$T = \begin{pmatrix} \cos \alpha & \sin \alpha \cos \beta & \sin \alpha \sin \beta \\ -\sin \alpha & \cos \alpha \cos \beta & \cos \alpha \sin \beta \\ 0 & -\sin \beta & \cos \beta \end{pmatrix}. \quad (4.1)$$

Then

$$T^{-1} = T^t \begin{pmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha \cos \beta & \cos \alpha \cos \beta & -\sin \beta \\ \sin \alpha \sin \beta & \cos \alpha \sin \beta & \cos \beta \end{pmatrix}. \quad (4.2)$$

Following section 3 of [32], we define the moving-frames:

$$\tilde{\mathbf{H}} = \begin{pmatrix} \mathcal{H}^1 \\ \mathcal{H}^2 \\ \mathcal{H}^3 \end{pmatrix} = T \begin{pmatrix} H^1 \\ H^2 \\ H^3 \end{pmatrix}, \quad \tilde{\mathbf{v}} = \begin{pmatrix} \mathcal{U} \\ \mathcal{V} \\ \mathcal{W} \end{pmatrix} = T \begin{pmatrix} u \\ v \\ w \end{pmatrix}, \quad \begin{pmatrix} \mathcal{X} \\ \mathcal{Y} \\ \mathcal{Z} \end{pmatrix} = T \begin{pmatrix} x \\ y \\ z \end{pmatrix}. \quad (4.3)$$

Write

$$\tilde{\nabla} = (\partial_{\mathcal{X}}, \partial_{\mathcal{Y}}, \partial_{\mathcal{Z}}) \quad \text{and} \quad \widetilde{\text{rot}} \mathbf{M} = \tilde{\nabla} \times \mathbf{M}. \quad (4.4)$$

Then we have

Lemma 4.1. *We have the following equations:*

$$\tilde{\mathbf{H}} \times \widetilde{\text{rot}} \tilde{\mathbf{H}} = T(\mathbf{H} \times \text{rot} \mathbf{H}), \quad (4.5)$$

$$\widetilde{\text{rot}}(\tilde{\mathbf{v}} \times \tilde{\mathbf{H}}) = T(\text{rot}(\mathbf{v} \times \mathbf{H})). \quad (4.6)$$

Proof. It follows from straightforward calculation. $\square$

We write

$$\begin{aligned} R_1 &= \mathcal{U}_t + \alpha'(\mathcal{Y}\mathcal{U}_{\mathcal{X}} - \mathcal{X}\mathcal{U}_{\mathcal{Y}} - \mathcal{V}) + (\mathcal{Z}\mathcal{U}_{\mathcal{X}} - \mathcal{X}\mathcal{U}_{\mathcal{Z}} - \mathcal{W})\beta' \sin \alpha \\ &\quad + (\mathcal{Z}\mathcal{U}_{\mathcal{Y}} - \mathcal{Y}\mathcal{U}_{\mathcal{Z}} - \mathcal{W})\beta' \cos \alpha + \mathcal{U}\mathcal{U}_{\mathcal{X}} + \mathcal{V}\mathcal{U}_{\mathcal{Y}} + \mathcal{W}\mathcal{U}_{\mathcal{Z}} - \nu \Delta(\mathcal{U}), \end{aligned} \quad (4.7)$$

$$\begin{aligned} R_2 &= \mathcal{V}_t + \alpha'(\mathcal{Y}\mathcal{V}_{\mathcal{X}} - \mathcal{X}\mathcal{V}_{\mathcal{Y}} + \mathcal{U}) + (\mathcal{Z}\mathcal{V}_{\mathcal{X}} - \mathcal{X}\mathcal{V}_{\mathcal{Z}})\beta' \sin \alpha \\ &\quad + (\mathcal{Z}\mathcal{V}_{\mathcal{Y}} - \mathcal{Y}\mathcal{V}_{\mathcal{Z}} - \mathcal{W})\beta' \cos \alpha + \mathcal{U}\mathcal{V}_{\mathcal{X}} + \mathcal{V}\mathcal{V}_{\mathcal{Y}} + \mathcal{W}\mathcal{V}_{\mathcal{Z}} - \nu \Delta(\mathcal{V}), \end{aligned} \quad (4.8)$$

$$\begin{aligned} R_3 &= \mathcal{W}_t + \alpha'(\mathcal{Y}\mathcal{W}_{\mathcal{X}} - \mathcal{X}\mathcal{W}_{\mathcal{Y}}) + (\mathcal{Z}\mathcal{W}_{\mathcal{X}} - \mathcal{X}\mathcal{W}_{\mathcal{Z}} + \mathcal{U})\beta' \sin \alpha \\ &\quad + (\mathcal{Z}\mathcal{W}_{\mathcal{Y}} - \mathcal{Y}\mathcal{W}_{\mathcal{Z}} + \mathcal{V})\beta' \cos \alpha + \mathcal{U}\mathcal{W}_{\mathcal{X}} + \mathcal{V}\mathcal{W}_{\mathcal{Y}} + \mathcal{W}\mathcal{W}_{\mathcal{Z}} - \nu \Delta(\mathcal{W}), \end{aligned} \quad (4.9)$$

$$\begin{cases} G_1 = \mathcal{H}^2(\mathcal{H}_{\mathcal{X}}^2 - \mathcal{H}_{\mathcal{Y}}^1) - \mathcal{H}^3(\mathcal{H}_{\mathcal{Z}}^1 - \mathcal{H}_{\mathcal{X}}^2), \\ G_2 = \mathcal{H}^3(\mathcal{H}_{\mathcal{Y}}^3 - \mathcal{H}_{\mathcal{Z}}^2) - \mathcal{H}^1(\mathcal{H}_{\mathcal{X}}^2 - \mathcal{H}_{\mathcal{Y}}^3), \\ G_3 = \mathcal{H}^1(\mathcal{H}_{\mathcal{Z}}^1 - \mathcal{H}_{\mathcal{X}}^3) - \mathcal{H}^2(\mathcal{H}_{\mathcal{Y}}^3 - \mathcal{H}_{\mathcal{Z}}^2), \end{cases} \quad (4.10)$$

$$\Phi_j = R_j - \mu_0 G_j \quad j = 1, 2, 3, \quad (4.11)$$

$$\begin{aligned} \Psi_1 &= \mathcal{H}_t^1 + \alpha'(\mathcal{Y}\mathcal{H}_{\mathcal{X}}^1 - \mathcal{X}\mathcal{H}_{\mathcal{Y}}^1 - \mathcal{H}^2) + (\mathcal{Z}\mathcal{H}_{\mathcal{X}}^1 - \mathcal{X}\mathcal{H}_{\mathcal{Z}}^1 - \mathcal{H}^3)\beta' \sin \alpha \\ &\quad + (\mathcal{Z}\mathcal{H}_{\mathcal{Y}}^1 - \mathcal{Y}\mathcal{H}_{\mathcal{Z}}^1)\beta' \cos \alpha - (\mathcal{U}\mathcal{H}^2 - \mathcal{V}\mathcal{H}^1)_{\mathcal{Y}} + (\mathcal{W}\mathcal{H}^1 - \mathcal{U}\mathcal{H}^3)_{\mathcal{Z}} - \eta \Delta \mathcal{H}^1, \end{aligned} \quad (4.12)$$

$$\begin{aligned}\Psi_2 &= \mathcal{H}_t^2 + \alpha'(\mathcal{Y}\mathcal{H}_\mathcal{X}^2 - \mathcal{X}\mathcal{H}_\mathcal{Y}^2 + \mathcal{H}^1) + (\mathcal{Z}\mathcal{H}_\mathcal{X}^2 - \mathcal{X}\mathcal{H}_\mathcal{Z}^2)\beta' \sin \alpha \\ &+ (\mathcal{Z}\mathcal{H}_\mathcal{Y}^2 - \mathcal{Y}\mathcal{H}_\mathcal{Z}^2 - \mathcal{H}^3)\beta' \cos \alpha - (\mathcal{V}\mathcal{H}^3 - \mathcal{W}\mathcal{H}^2)_\mathcal{Z} + (\mathcal{U}\mathcal{H}^2 - \mathcal{V}\mathcal{H}^1)_\mathcal{X} - \eta\Delta\mathcal{H}^2,\end{aligned}\quad (4.13)$$

$$\begin{aligned}\Psi_3 &= \mathcal{H}_t^3 + \alpha'(\mathcal{Y}\mathcal{H}_\mathcal{X}^3 - \mathcal{X}\mathcal{H}_\mathcal{Y}^3) + (\mathcal{Z}\mathcal{H}_\mathcal{X}^3 - \mathcal{X}\mathcal{H}_\mathcal{Z}^3 + \mathcal{H}^1)\beta' \sin \alpha \\ &+ (\mathcal{Z}\mathcal{H}_\mathcal{Y}^3 - \mathcal{Y}\mathcal{H}_\mathcal{Z}^3 + \mathcal{H}^2)\beta' \cos \alpha - (\mathcal{W}\mathcal{H}^1 - \mathcal{U}\mathcal{H}^3)_\mathcal{X} + (\mathcal{V}\mathcal{H}^3 - \mathcal{W}\mathcal{H}^2)_\mathcal{Y} - \eta\Delta\mathcal{H}^3.\end{aligned}\quad (4.14)$$

Then the MHD equations (1.1)-(1.4) become

$$\Phi_1 + \frac{1}{\rho}p_\mathcal{X} = 0, \quad \Phi_2 + \frac{1}{\rho}p_\mathcal{Y} = 0, \quad \Phi_3 + \frac{1}{\rho}p_\mathcal{Z} = 0, \quad (4.15)$$

$$\Psi_j = 0, \quad j = 1, 2, 3, \quad (4.16)$$

$$\begin{cases} \mathcal{U}_\mathcal{X} + \mathcal{V}_\mathcal{Y} + \mathcal{W}_\mathcal{Z} = 0, \\ \mathcal{H}_\mathcal{X}^1 + \mathcal{H}_\mathcal{Y}^2 + \mathcal{H}_\mathcal{Z}^3 = 0. \end{cases} \quad (4.17)$$

Instead of solving (4.15), we will first deal with the compatibility conditions:

$$\partial_\mathcal{Y}(\Phi_1) = \partial_\mathcal{X}(\Phi_2), \quad \partial_\mathcal{Z}(\Phi_1) = \partial_\mathcal{X}(\Phi_3), \quad \partial_\mathcal{Z}(\Phi_2) = \partial_\mathcal{Y}(\Phi_3). \quad (4.18)$$

Then we find p via (4.15).

Now we suppose that

$$\begin{cases} \mathcal{U} = -2\gamma\mathcal{X} - \alpha\mathcal{Y} - \mathcal{Z}\beta' \sin \alpha, \\ \mathcal{V} = \phi + \gamma\mathcal{Y}, \\ \mathcal{W} = \psi + \gamma\mathcal{Z}, \end{cases} \quad \text{and} \quad \begin{cases} \mathcal{H}^1 = g - 2\xi\mathcal{X}, \\ \mathcal{H}^2 = \sigma + \xi\mathcal{Y}, \\ \mathcal{H}^3 = h + \xi\mathcal{Z}, \end{cases} \quad (4.19)$$

where ϕ, ψ, σ and h are functions in $\mathcal{X}$ and t , and g, γ and ξ are functions in t .

Substituting (4.19) into (4.16)-(4.18), we get that

$$\begin{cases} \xi' = 0, \\ g' = -2\gamma g \end{cases} \quad (4.20)$$

and

$$\begin{cases} \phi_t - 2\gamma\mathcal{X}\phi_\mathcal{X} + \gamma\phi - \nu\phi_\mathcal{X}\mathcal{X} - \psi\beta' \cos \alpha + \mu_0(\xi\sigma + g\sigma_\mathcal{X} - 2\mathcal{X}\xi\sigma_\mathcal{X}) \\ \quad = ((\beta')^2 \sin \alpha \cos \alpha + \alpha'\gamma - \alpha'')\mathcal{X} + \theta_1, \\ \psi_\mathcal{Y} - 2\gamma\mathcal{X}\psi_\mathcal{X} + \gamma\psi - \nu\psi_\mathcal{X}\mathcal{X} + \phi\beta' \cos \alpha + \mu_0(\xi h + gh_\mathcal{X} - 2\mathcal{X}\xi h_\mathcal{X}) \\ \quad = -((\beta' \sin \alpha)' + \alpha'\beta' \cos \alpha + \alpha'\beta' \cos \alpha - \gamma'\beta' \sin \alpha)\mathcal{X} + \theta_2, \\ \sigma_t - 2\gamma\mathcal{X}\sigma_\mathcal{X} - \gamma\sigma - \eta\sigma_\mathcal{X}\mathcal{X} - h\beta' \cos \alpha + (\xi\phi - g\phi_\mathcal{X} + 2\mathcal{X}\xi\phi_\mathcal{X}) \\ \quad = 3\alpha'\mathcal{X}\xi - \alpha'g, \\ \mathcal{H}_t - 2\gamma\mathcal{X}h_\mathcal{X} - \gamma h - \eta h_\mathcal{X}\mathcal{X} + \sigma\beta' \cos \alpha + (\xi\psi - g\psi_\mathcal{X} + 2\mathcal{X}\xi\psi_\mathcal{X}) \\ \quad = 3\mathcal{X}\beta'\xi \sin \alpha - g\beta' \sin \alpha, \end{cases} \quad (4.21)$$

where θ_1 and θ_2 are any functions of t .

Set

$$\varphi = \int \beta' \cos \alpha dt \quad \text{and} \quad \mu = \exp(4 \int \gamma dt). \quad (4.22)$$

We have $g = c/\sqrt{\mu}$ by (4.20), where c is a constant.

Change variables by

$$\begin{pmatrix} \phi \\ \psi \\ \sigma \\ h \end{pmatrix} = \begin{pmatrix} \mu^{-\frac{1}{4}} \cos \varphi & \mu^{-\frac{1}{4}} \sin \varphi & 0 & 0 \\ -\mu^{-\frac{1}{4}} \sin \varphi & \mu^{-\frac{1}{4}} \cos \varphi & 0 & 0 \\ 0 & 0 & \mu^{\frac{1}{4}} \cos \varphi & \mu^{\frac{1}{4}} \sin \varphi \\ 0 & 0 & -\mu^{\frac{1}{4}} \sin \varphi & \mu^{\frac{1}{4}} \cos \varphi \end{pmatrix} \begin{pmatrix} \hat{\phi} \\ \hat{\psi} \\ \hat{\sigma} \\ \hat{h} \end{pmatrix}. \quad (4.23)$$

Then (4.21) becomes

$$\begin{cases} \hat{\phi}_t - \frac{\mu'}{2\mu} \mathcal{X} \hat{\phi}_{\mathcal{X}} - \nu \hat{\phi}_{\mathcal{X}\mathcal{X}} + \mu_0 \sqrt{\mu} (\xi \hat{\sigma} + \frac{c}{\sqrt{\mu}} \hat{\sigma}_{\mathcal{X}} - 2\xi \mathcal{X} \hat{\sigma}_{\mathcal{X}}) = q_1 \mathcal{X} + \beta_1, \\ \hat{\sigma}_t - \frac{\mu'}{2\mu} \mathcal{X} \hat{\sigma}_{\mathcal{X}} - \eta \hat{\sigma}_{\mathcal{X}\mathcal{X}} + \frac{\xi}{\sqrt{\mu}} \hat{\phi} - \frac{c}{\mu} \hat{\phi}_{\mathcal{X}} + 2\frac{\xi}{\sqrt{\mu}} \mathcal{X} \hat{\phi}_{\mathcal{X}} = q_2 \mathcal{X} + \beta_2, \\ \hat{\psi}_t - \frac{\mu'}{2\mu} \mathcal{X} \hat{\psi}_{\mathcal{X}} - \nu \hat{\psi}_{\mathcal{X}\mathcal{X}} + \mu_0 \sqrt{\mu} (\xi \hat{h} + \frac{c}{\sqrt{\mu}} \hat{h}_{\mathcal{X}} - 2\xi \mathcal{X} \hat{h}_{\mathcal{X}}) = q_3 \mathcal{X} + \beta_3, \\ \hat{h}_t - \frac{\mu'}{2\mu} \mathcal{X} \hat{h}_{\mathcal{X}} - \eta \hat{h}_{\mathcal{X}\mathcal{X}} + \frac{\xi}{\sqrt{\mu}} \hat{\psi} - \frac{c}{\mu} \hat{\psi}_{\mathcal{X}} + \frac{2\xi}{\sqrt{\mu}} \mathcal{X} \hat{\psi}_{\mathcal{X}} = q_4 \mathcal{X} + \beta_4, \end{cases} \quad (4.24)$$

where

$$\begin{cases} q_1 = \mu^{\frac{1}{4}} (((\beta')^2 \sin \alpha \cos \alpha + \alpha' \gamma - \alpha'') \cos \varphi - ((\beta' \sin \alpha)' + (\alpha' - \gamma') \beta' \cos \alpha) \sin \varphi), \\ q_2 = \mu^{\frac{1}{4}} (-((\beta')^2 \sin \alpha \cos \alpha + \alpha' \gamma - \alpha'') \sin \varphi - ((\beta' \sin \alpha)' + (\alpha' - \gamma') \beta' \cos \alpha) \cos \varphi), \\ q_3 = \mu^{-\frac{1}{4}} (3\alpha' \xi \cos \varphi + 3\beta' \xi \sin \alpha \sin \varphi), \\ q_4 = \mu^{-\frac{1}{4}} (-3\alpha' \xi \sin \varphi + 3\beta' \xi \sin \alpha \cos \varphi) \end{cases} \quad (4.25)$$

and

$$\begin{cases} \beta_1 = \mu^{\frac{1}{4}} (\theta_1 \cos \varphi - \theta_2 \sin \varphi), \\ \beta_2 = \mu^{\frac{1}{4}} (-\theta_1 \sin \varphi - \theta_2 \cos \varphi), \\ \beta_3 = \mu^{-\frac{1}{4}} (-\alpha' g \cos \varphi - \beta' \xi \sin \alpha \sin \varphi), \\ \beta_4 = \mu^{-\frac{1}{4}} (\alpha' g \sin \varphi - \beta' \xi \sin \alpha \cos \varphi). \end{cases} \quad (4.26)$$

Let n be a positive integer. We write

$$\hat{\phi} = \sum_{m=0}^n a_m(t) \mathcal{X}^m, \quad \hat{\sigma} = \sum_{m=0}^n b_m(t) \mathcal{X}^m, \quad \hat{\psi} = \sum_{m=0}^n c_m(t) \mathcal{X}^m, \quad \hat{h} = \sum_{m=0}^n d_m(t) \mathcal{X}^m. \quad (4.27)$$

Substituting above expressions into (4.24), we obtain that

$$\begin{cases} a_{m,t} - (m+1) \frac{\mu'}{\mu} a_m - \xi \mu_0 \sqrt{\mu} (2m+1) b_m = A_m(t), \\ b_{m,t} - (m+1) \frac{\mu'}{2\mu} b_m + (2m+3) \frac{\xi}{\sqrt{\mu}} a_m = B_m(t), \end{cases} \quad (4.28)$$

where

$$\begin{cases} A_m = -c \mu_0 (m+1) b_{m+1} + (m+1)(m+2) \nu a_{m+2} + \delta_{m,1} q_1 + \delta_{m,0} \beta_1, \\ B_m = \frac{c}{\mu} (m+1) a_{m+1} + (m+1)(m+2) \eta b_{m+2} + \delta_{m,1} q_2 + \delta_{m,0} \beta_2 \end{cases} \quad (4.29)$$

and

$$\begin{cases} c_{m,t} - (m+1) \frac{\mu'}{\mu} c_m - \xi \mu_0 \sqrt{\mu} (2m+1) d_m = C_m(t), \\ d_{m,t} - (m+1) \frac{\mu'}{2\mu} d_m + (2m+3) \frac{\xi}{\sqrt{\mu}} c_m = D_m(t), \end{cases} \quad (4.30)$$

where

$$\begin{cases} C_m = -c\mu_0(m+1)d_{m+1} + (m+1)(m+2)\nu c_{m+2} + \delta_{m,1}q_3 + \delta_{m,0}\beta_3, \\ D_m = \frac{c}{\mu}(m+1)c_{m+1} + (m+1)(m+2)\eta d_{m+2} + \delta_{m,1}q_4 + \delta_{m,0}\beta_4. \end{cases} \quad (4.31)$$

As we did before, let

$$\mu = l_1 \exp\left(\int (2k + \frac{4}{-1 + 2l_2 e^{-kt}}) dt\right) \quad (4.32)$$

for any constants k , l_1 and l_2 . Under this condition, the equations (4.28) and (4.30) may be solved as second order constant-coefficient ordinary differential equations via certain changing of variables. We only give the results but omit the procedures.

Write $\lambda_m = \frac{1}{4}k^2 - (2m+1)(2m+3)\xi^2\mu_0$ and define

$$\zeta_{m,1} = \begin{cases} \mu \exp(\sqrt{\lambda_m} t) & \text{if } \lambda_m > 0, \\ \mu & \text{if } \lambda_m = 0, \\ \mu \cos(i\sqrt{\lambda_m} t) & \text{if } \lambda_m < 0, \end{cases} \quad (4.33)$$

$$\zeta_{m,2} = \begin{cases} \mu \exp(-\sqrt{\lambda_m} t) & \text{if } \lambda_m > 0, \\ t\mu & \text{if } \lambda_m = 0, \\ \mu \sin(i\sqrt{\lambda_m} t) & \text{if } \lambda_m < 0. \end{cases} \quad (4.34)$$

In addition, we denote

$$\begin{cases} P_m = A'_m - A_m - \frac{\mu'}{\mu}(m+1)A_m + \xi\mu_0(2m+1)\sqrt{\mu}B_m, \\ Q_m = C'_m - C_m - \frac{\mu'}{\mu}(m+1)C_m + \xi\mu_0(2m+1)\sqrt{\mu}D_m. \end{cases} \quad (4.35)$$

Then we get the following recursive formulae:

$$a_m = \sum_{j=1}^2 \varsigma_{m,j}(q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} P_m(s) ds), \quad (4.36)$$

$$\begin{aligned} b_m &= \frac{1}{(2m+1)\xi\mu_0\sqrt{\mu}} \sum_{j=1}^2 \left(\varsigma'_{m,j}(q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} P_m(s) ds \right. \\ &+ \left. \varsigma_{m,j} \left(\frac{W_j[\varsigma_{m,1}(t), \varsigma_{m,2}(t)]}{W[\varsigma_{m,1}(t), \varsigma_{m,2}(t)]} P_m(t) - (m+1) \frac{\mu'}{2\mu} (q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} P_m(s) ds) \right) \right), \end{aligned} \quad (4.37)$$

$$c_m = \sum_{j=1}^2 \varsigma_{m,j}(q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} Q_m(s) ds), \quad (4.38)$$

and

$$\begin{aligned} d_m &= \frac{1}{(2m+1)\xi\mu_0\sqrt{\mu}} \sum_{j=1}^2 \left(\varsigma'_{m,j}(q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} Q_m(s) ds \right. \\ &+ \left. \varsigma_{m,j} \left(\frac{W_j[\varsigma_{m,1}(t), \varsigma_{m,2}(t)]}{W[\varsigma_{m,1}(t), \varsigma_{m,2}(t)]} Q_m(t) - (m+1) \frac{\mu'}{2\mu} (q_{m,j} + \int_0^t \frac{W_j[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]}{W[\varsigma_{m,1}(s), \varsigma_{m,2}(s)]} Q_m(s) ds) \right) \right), \end{aligned} \quad (4.39)$$

where $q_{m,1}$ and $q_{m,2}$ are real constants.

Theorem 4.2. Let α and β be arbitrary functions of t , and let c and ξ be arbitrary real constants. The functions φ , μ , $\hat{\phi}$, $\hat{\sigma}$, $\hat{\psi}$ and $\hat{h}$ are given in (4.22), (4.32), (4.27) and (4.36)-(4.39), respectively. Then we have the following solutions of the MHD equations (1.1)-(1.4):

$$\left\{ \begin{array}{l} u = -\mathcal{X} \frac{2\mu'}{\mu} \cos \alpha - \mathcal{Y}(\alpha' \cos \alpha + \frac{\mu'}{4\mu} \sin \alpha) - \mathcal{Z}\beta' \sin \alpha \cos \alpha - \mu^{-\frac{1}{4}}(\hat{\phi} \cos \varphi - \hat{\psi} \sin \varphi) \sin \alpha, \\ v = -\mathcal{X} \frac{2\mu'}{\mu} \sin \alpha \cos \beta + \mathcal{Y}(-\alpha' \sin \alpha + \frac{\mu'}{4\mu} \cos \alpha) \cos \beta + \mathcal{Z}(-\beta' \sin \alpha^2 \cos \beta + \frac{\mu'}{4\mu} \sin \beta) \\ \quad + \mu^{-\frac{1}{4}}((\cos \alpha \cos \beta \cos \varphi + \sin \beta \sin \varphi)\hat{\phi} + (\cos \alpha \cos \beta \sin \varphi - \sin \beta \cos \varphi)\hat{\psi}), \\ w = -\mathcal{X} \frac{2\mu'}{\mu} \sin \alpha \sin \beta + \mathcal{Y}(-\alpha' \sin \alpha + \frac{\mu'}{4\mu} \cos \alpha) \sin \beta + \mathcal{Z}(-\beta' \sin \alpha^2 \sin \beta + \frac{\mu'}{4\mu} \cos \beta) \\ \quad + \mu^{-\frac{1}{4}}((\cos \alpha \sin \beta \cos \varphi - \cos \beta \sin \varphi)\hat{\phi} + (\cos \alpha \sin \beta \sin \varphi + \cos \beta \cos \varphi)\hat{\psi}), \end{array} \right. \quad (4.40)$$

$$\left\{ \begin{array}{l} H^1 = \frac{c}{\sqrt{\mu}} \cos \alpha - 2\mathcal{X}\xi \cos \alpha - \mathcal{Y}\xi \sin \alpha - \mu^{\frac{1}{4}} \sin \alpha (\hat{\sigma} \cos \varphi + \hat{h} \sin \varphi), \\ H^2 = \frac{c}{\sqrt{\mu}} \sin \alpha \cos \beta - 2\mathcal{X}\xi \sin \alpha \cos \beta + \mathcal{Y}\xi \cos \alpha \cos \beta - \mathcal{Z}\xi \sin \beta \\ \quad + \mu^{-\frac{1}{4}}((\cos \alpha \cos \beta \cos \varphi + \sin \beta \sin \varphi)\hat{\sigma} + (\cos \alpha \cos \beta \sin \varphi - \sin \beta \cos \varphi)\hat{h}), \\ H^3 = \frac{c}{\sqrt{\mu}} \sin \alpha \sin \beta - 2\mathcal{X}\xi \sin \alpha \sin \beta + \mathcal{Y}\xi \cos \alpha \cos \beta + \mathcal{Z}\xi \cos \beta \\ \quad + \mu^{-\frac{1}{4}}((\cos \alpha \sin \beta \cos \varphi - \cos \beta \sin \varphi)\hat{\sigma} + (\cos \alpha \sin \beta \sin \varphi + \cos \beta \cos \varphi)\hat{h}), \end{array} \right. \quad (4.41)$$

$$\begin{aligned} p = & -\rho \left(\frac{-2\mu^2\mu'' - (\mu')^3 + \mu\mu'\mu'' + 2\mu(\mu')^2}{4\mu^3} + (\alpha')^2 + (\beta' \sin \alpha)^2 \right) \frac{\mathcal{X}^2}{2} + \left(\frac{(\mu')^2}{16\mu^2} - (\alpha')^2 \right) \frac{\mathcal{Y}^2}{2} \\ & + \left(\frac{(\mu')^2}{16\mu^2} - (\beta' \sin \alpha)^2 \right) \frac{\mathcal{Z}^2}{2} - (\alpha'' + \frac{\mu'}{2\mu} \alpha' - \beta' \sin \alpha \beta' \cos \alpha) \mathcal{X}\mathcal{Y} \\ & - ((\beta' \sin \alpha)' + \alpha' \beta' \cos \alpha + \frac{\mu'}{2\mu} \beta' \sin \alpha) \mathcal{X}\mathcal{Z} - \mathcal{Y}\mathcal{Z} \alpha' \beta' \sin \alpha - 2\alpha' \int \phi d\mathcal{X} \\ & - 2\beta' \sin \alpha \int \psi d\mathcal{X} - \mu_0 \left(\frac{1}{2} (\sigma^2 + h^2) + \xi \sigma \mathcal{Y} + \xi, h \mathcal{Z} \right) \end{aligned} \quad (4.42)$$

where ϕ , ψ , σ and h are given by (4.23).

Set

$$\left\{ \begin{array}{l} \mathcal{U} = a_{11}\mathcal{X} + a_{12}\mathcal{Y} + a_{13}\mathcal{Z} - 6\nu\mathcal{X}^{-1}, \\ \mathcal{V} = a_{21}\mathcal{X} + a_{22}\mathcal{Y} + a_{23}\mathcal{Z} - 6\nu\mathcal{Y}\mathcal{X}^{-2}, \\ \mathcal{W} = a_{31}\mathcal{X} + a_{32}\mathcal{Y} + a_{33}\mathcal{Z}, \end{array} \right. \quad \left\{ \begin{array}{l} \mathcal{H}^1 = b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z}, \\ \mathcal{H}^2 = b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z}, \\ \mathcal{H}^3 = b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}. \end{array} \right. \quad (4.43)$$

Substituting (4.43) into (4.18), one gets that

$$a_{11} = a_{22} = \gamma, \quad a_{21} = -a_{12} = \alpha', \quad a_{31} = -a_{13} = \beta' \sin \alpha, \quad a_{23} = a_{32} = -\beta' \cos \alpha \quad (4.44)$$

and

$$\left\{ \begin{array}{l} 2\alpha'' + 4\alpha'\gamma = \mu_0(b_{31}(b_{32} - b_{23}) + b_{33}(b_{21} - b_{12}) + b_{32}(b_{13} - b_{31})), \\ b_{22}(b_{13} - b_{31}) = -b_{23}(b_{21} - b_{12}) - b_{21}(b_{32} - b_{23}), \\ b_{11}(b_{32} - b_{23}) = b_{12}(b_{32} - b_{23}) - b_{13}(b_{21} - b_{12}). \end{array} \right. \quad (4.45)$$

By (4.16), we have that

$$\begin{aligned} & b'_{11}\mathcal{X} + b'_{12}\mathcal{Y} + b'_{13}\mathcal{Z} + b_{11}(\alpha'\mathcal{Y} + \mathcal{Z}\beta' \sin \alpha) + b_{12}(\gamma\mathcal{Y} - 6\nu\mathcal{Y}\mathcal{X}^{-2}) \\ & - 2(\mathcal{X}\beta' \sin \alpha + \mathcal{Y}\beta' \cos \alpha + \gamma\mathcal{Z})b_{13} - (\gamma + 6\nu\mathcal{X}^{-2})(b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z}) \\ & + b_{11}(\gamma\mathcal{X} - \alpha'\mathcal{Y} - \mathcal{Z}\beta' \sin \alpha - 6\nu\mathcal{X}^{-1}) = 0, \end{aligned} \quad (4.46)$$

$$\begin{aligned}
& b'_{21}\mathcal{X} + b'_{22}\mathcal{Y} + b'_{23}\mathcal{Z} + b_{21}(\gamma\mathcal{X} - 6\nu\mathcal{X}^{-1}) + b_{22}(-\alpha'\mathcal{X} + \mathcal{Z}\beta'\cos\alpha) \\
& - 2(\mathcal{X}\beta'\sin\alpha + \mathcal{Y}\beta'\cos\alpha + \gamma\mathcal{Z})b_{23} + (\gamma + 6\nu\mathcal{X}^{-2})(b_{12}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z}) \\
& + b_{22}(\alpha'\mathcal{X} + \gamma\mathcal{Y} - \mathcal{Z}\beta'\cos\alpha - 6\nu\mathcal{Y}\mathcal{X}^{-2}) = 0
\end{aligned} \tag{4.47}$$

and

$$\begin{aligned}
& b'_{31}\mathcal{X} + b'_{32}\mathcal{Y} + b'_{33}\mathcal{Z} + b_{31}(\gamma\mathcal{X} - 6\nu\mathcal{X}^{-1}) + b_{32}(\gamma\mathcal{Y} - 6\nu\mathcal{Y}\mathcal{X}^{-2}) \\
& - b_{33}(\mathcal{X}\beta'\sin\alpha + \mathcal{Y}\beta'\cos\alpha) + 2(b_{21}\mathcal{X} + b_{22}\mathcal{Y})\beta'\cos\alpha \\
& + 2\gamma(b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}) - b_{33}(\mathcal{X}\beta'\sin\alpha + \mathcal{Y}\beta'\cos\alpha + 2\gamma\mathcal{Z}) = 0.
\end{aligned} \tag{4.48}$$

One shows that $\tilde{\mathbf{H}} \neq 0$ if and only if β is a constant. In this case,

$$b_{11} = b_{12} = b_{13} = b_{23} = b_{31} = b_{32} = 0 \tag{4.49}$$

and

$$b'_{22} = b'_{21} = 0. \tag{4.50}$$

Hence, one gets

$$\gamma = -\frac{\alpha''}{2\alpha'} - \frac{\mu_0 b_{22} b_{21}}{4\alpha'}. \tag{4.51}$$

Theorem 4.3. *For any function $\alpha(t)$ and constants a, b , we have a solution of the MHD equations (1.1)-(1.4):*

$$\begin{cases} u = -\left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'}\right)x - \alpha'y - \frac{6\nu \cos\alpha}{x \cos\alpha + y \sin\alpha} + 6\nu \sin\alpha \frac{-x \sin\alpha + y \cos\alpha}{(x \cos\alpha + y \sin\alpha)^2}, \\ v = \alpha'x - \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'}\right)y - \frac{6\nu \sin\alpha}{x \cos\alpha + y \sin\alpha} - 6\nu \cos\alpha \frac{-x \sin\alpha + y \cos\alpha}{(x \cos\alpha + y \sin\alpha)^2}, \\ w = 2\left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'}\right)z, \end{cases} \tag{4.52}$$

$$\begin{cases} H^1 = -[(a \cos\alpha - b \sin\alpha)x + (-a \sin\alpha + b \cos\alpha)y] \sin\alpha, \\ H^2 = [(a \cos\alpha - b \sin\alpha)x + (a \sin\alpha + b \cos\alpha)y] \cos\alpha, \\ H^3 = -bz \end{cases} \tag{4.53}$$

and

$$\begin{aligned}
p = & -\rho \left(\left(\left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)' + \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)^2 - (\alpha')^2 - \mu_0 a^2 \right) \frac{(x \cos\alpha + y \sin\alpha)^2}{2} \right. \\
& + \left(\left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)' + \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)^2 - (\alpha')^2 \right) \frac{(-x \sin\alpha + y \cos\alpha)^2}{2} \\
& - \left(\left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)' + 2 \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)^2 \right) \frac{z^2}{2} \\
& + (\alpha'' + 2\alpha' \left(\frac{\alpha''}{2\alpha'} + \frac{\mu_0 ab}{4\alpha'} \right)) (x \cos\alpha + y \sin\alpha) (-x \sin\alpha + y \cos\alpha) \\
& \left. - 12\nu\alpha' \frac{-x \sin\alpha + y \cos\alpha}{x \cos\alpha + y \sin\alpha} + \frac{12\nu^2}{(x \cos\alpha + y \sin\alpha)^2} \right).
\end{aligned} \tag{4.54}$$

Let α_1, β_1 and γ be functions of t and a, b be real constants. Set

$$\xi_r = \begin{cases} e^{\alpha_1\mathcal{Y}+\beta_1\mathcal{Z}} - ae^{-(\alpha_1\mathcal{Y}+\beta_1\mathcal{Z})} & \text{if } r = 0, \\ \sin(\alpha_1\mathcal{Y} + \beta_1\mathcal{Z}) & \text{if } r = 1; \end{cases} \tag{4.55}$$

$$\zeta_r = \begin{cases} e^{\alpha_1\mathcal{Y}+\beta_1\mathcal{Z}} + ae^{-(\alpha_1\mathcal{Y}+\beta_1\mathcal{Z})} & \text{if } r = 0, \\ \cos(\alpha_1\mathcal{Y} + \beta_1\mathcal{Z}) & \text{if } r = 1; \end{cases} \tag{4.56}$$

$$\phi_s = \begin{cases} e^{\gamma\mathcal{X}} - be^{-\gamma\mathcal{X}} & \text{if } s = 0, \\ \sin(\gamma\mathcal{X}) & \text{if } s = 1 \end{cases} \quad \text{and } \psi_s = \begin{cases} e^{\gamma\mathcal{X}} + be^{-\gamma\mathcal{X}} & \text{if } s = 0, \\ \cos(\gamma\mathcal{X}) & \text{if } s = 1. \end{cases} \quad (4.57)$$

We assume that

$$\begin{pmatrix} \mathcal{U} \\ \mathcal{V} \\ \mathcal{W} \end{pmatrix} = A \begin{pmatrix} \mathcal{X} \\ \mathcal{Y} \\ \mathcal{Z} \end{pmatrix} + \begin{pmatrix} -(\alpha_1\sigma + \beta_1\tau)\zeta_r\phi_s \\ \sigma\gamma\xi_r\psi_s \\ \tau\gamma\xi_r\psi_s \end{pmatrix}, \quad (4.58)$$

$$\begin{pmatrix} \mathcal{H}^1 \\ \mathcal{H}^2 \\ \mathcal{H}^3 \end{pmatrix} = B \begin{pmatrix} \mathcal{X} \\ \mathcal{Y} \\ \mathcal{Z} \end{pmatrix} + \begin{pmatrix} -(\alpha_1\sigma_1 + \beta_1\tau_1)\zeta_r\phi_s \\ \sigma_1\gamma\xi_r\psi_s \\ \tau_1\gamma\xi_r\psi_s \end{pmatrix}, \quad (4.59)$$

where $A = (a_{i,j})_{3 \times 3}$ and $B = (b_{i,j})_{3 \times 3}$ are 3×3 matrices whose entries are functions of t ; the functions σ , τ , σ_1 and τ_1 are also functions of t ; the numbers r and s may be 0 or 1.

Now

$$\begin{aligned} R_1 = & (a'_{11} + a_{11}^2 + a_{12}(a_{21} - \alpha') + a_{13}(a_{31} - \beta' \sin \alpha) - a_{21}\alpha' - a_{31}\beta' \sin \alpha)\mathcal{X} \\ & + (a'_{12} + a_{11}(a_{12} + \alpha') + a_{12}a_{22} + a_{13}(a_{32} - \beta' \cos \alpha) - a_{22}\alpha' - a_{32}\beta' \sin \alpha)\mathcal{Y} \\ & + (a'_{13} + a_{11}(a_{13} + \beta' \sin \alpha) + a_{12}(a_{23} + \beta' \cos \alpha) + a_{13}a_{33} - a_{23}\alpha' - a_{33}\beta' \sin \alpha)\mathcal{Z} \\ & + (\delta_{r,1} + 4a\delta_{r,0})(\alpha_1\sigma + \beta_1\tau)^2\gamma\phi_s\psi_s \\ & - (-1)^r(\alpha_1\sigma + \beta_1\tau)[\alpha'_1\mathcal{Y} + \beta'_1\mathcal{Z} + \alpha_1((a_{21} - \alpha')\mathcal{X} + a_{22}\mathcal{Y} + (a_{23} + \beta' \cos \alpha)\mathcal{Z}) \\ & + \beta_1((a_{31} - \beta' \sin \alpha)\mathcal{X} + (a_{32} - \beta' \cos \alpha)\mathcal{Y} + a_{33}\mathcal{Z})]\xi_r\phi_s \\ & + \gamma[(a_{12} - \alpha')\sigma + (a_{13} - \beta' \sin \alpha)\tau]\xi_r\psi_s \\ & - [(\alpha_1\sigma + \beta_1\tau)' + a_{11}(\alpha_1\sigma + \beta_1\tau) + \nu(\alpha_1\sigma + \beta_1\tau)((-1)^s\gamma^2 + (-1)^r(\alpha_1^2 + \beta_1^2))]\zeta_r\phi_s \\ & - (\alpha_1\sigma + \beta_1\tau)[\gamma'\mathcal{X} + \gamma(a_{11}\mathcal{X} + (a_{12} + \alpha')\mathcal{Y} + (a_{13} + \beta' \sin \alpha)\mathcal{Z})]\zeta_r\psi_s, \end{aligned} \quad (4.60)$$

$$\begin{aligned} R_2 = & (a'_{21} + a_{21}a_{11} + a_{22}(a_{21} - \alpha') + a_{23}(a_{31} - \beta' \sin \alpha) + a_{11}\alpha' - a_{31}\beta' \cos \alpha)\mathcal{X} \\ & + (a'_{22} + a_{21}(a_{12} + \alpha') + a_{22}^2 + a_{23}(a_{32} - \beta' \cos \alpha) + a_{12}\alpha' - a_{32}\beta' \cos \alpha)\mathcal{Y} \\ & + (a'_{23} + a_{21}(a_{13} + \beta' \sin \alpha) + a_{22}(a_{23} + \beta' \cos \alpha) + a_{23}a_{33} + a_{13}\alpha' - a_{33}\beta' \cos \alpha)\mathcal{Z} \\ & + (\delta_{r,1} + 4a\delta_{r,0})\sigma\gamma^2(\alpha_1\sigma + \beta_1\tau)\xi_r\zeta_r \\ & + (-1)^s\sigma\gamma[\gamma'\mathcal{X} + \gamma(a_{11}\mathcal{X} + (a_{12} + \alpha')\mathcal{Y} + (a_{13} + \beta' \sin \alpha)\mathcal{Z})]\xi_r\phi_s \\ & + [(\sigma\gamma)' + a_{22}\sigma\gamma + (a_{23} - \beta' \cos \alpha)\tau\gamma - \nu\sigma\gamma((-1)^s\gamma^2 + (-1)^r(\alpha_1^2 + \beta_1^2))]\xi_r\psi_s \\ & - (a_{21} + \alpha')(\alpha_1\sigma + \beta_1\tau)\zeta_r\phi_s \\ & + \sigma\gamma[(\alpha'_1\mathcal{Y} + \beta'_1\mathcal{Z}) + \alpha_1((a_{21} - \alpha')\mathcal{X} + a_{22}\mathcal{Y} + (a_{23} + \beta' \cos \alpha)\mathcal{Z}) \\ & + \beta_1((a_{31} - \beta' \sin \alpha)\mathcal{X} + (a_{32} - \beta' \cos \alpha)\mathcal{Y} + a_{33}\mathcal{Z})]\zeta_r\psi_s, \end{aligned} \quad (4.61)$$

$$\begin{aligned} R_3 = & (a'_{31} + a_{31}a_{11} + a_{32}(a_{21} - \alpha') + a_{33}(a_{31} - \beta' \sin \alpha) + a_{11} + a_{21}\beta' \cos \alpha)\mathcal{X} \\ & + (a'_{32} + a_{31}(a_{12} + \alpha') + a_{32}a_{22} + a_{33}(a_{32} - \beta' \cos \alpha) + a_{12} + a_{22}\beta' \cos \alpha)\mathcal{Y} \\ & + (a'_{33} + a_{31}(a_{13} + \beta' \sin \alpha) + a_{32}(a_{23} + \beta' \cos \alpha) + a_{33}^2 + a_{13}\beta' \sin \alpha + a_{23}\beta' \cos \alpha)\mathcal{Z} \\ & + (\delta_{r,1} + 4a\delta_{r,0})\tau\gamma^2(\alpha_1\sigma + \beta_1\tau)\xi_r\zeta_r \\ & + (-1)^s\tau\gamma[\gamma'\mathcal{X} + \gamma(a_{11}\mathcal{X} + (a_{12} + \alpha')\mathcal{Y} + (a_{13} + \beta' \sin \alpha)\mathcal{Z})]\xi_r\phi_s \\ & + [(\tau\gamma)' + (a_{32} + \beta' \cos \alpha)\sigma\gamma + a_{33}\tau\gamma - \nu\tau\gamma((-1)^s\gamma^2 + (-1)^r(\alpha_1^2 + \beta_1^2))]\xi_r\psi_s \\ & - (a_{31} + \beta' \sin \alpha)(\alpha_1\sigma + \beta_1\tau)\zeta_r\phi_s \\ & + \tau\gamma[(\alpha'_1\mathcal{Y} + \beta'_1\mathcal{Z}) + \alpha_1((a_{21} - \alpha')\mathcal{X} + a_{22}\mathcal{Y} + (a_{23} + \beta' \cos \alpha)\mathcal{Z}) \\ & + \beta_1((a_{31} - \beta' \sin \alpha)\mathcal{X} + (a_{32} - \beta' \cos \alpha)\mathcal{Y} + a_{33}\mathcal{Z})]\zeta_r\psi_s, \end{aligned} \quad (4.62)$$

$$\begin{aligned} G_1 = & (b_{21} - b_{12})(b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z}) + (b_{31} - b_{13})(b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}) \\ & + \gamma[(-1)^s\gamma^2(\sigma_1^2 + \tau_1^2) + (-1)^r(\alpha_1\sigma_1 + \beta_1\tau_1)^2]\xi_r^2\phi_s\psi_s \\ & + [(b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z})((-1)^s\sigma_1\gamma^2 + (-1)^r\alpha_1(\alpha_1\sigma_1 + \beta_1\tau_1)) \\ & + (b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z})((-1)^s\tau_1\gamma^2 + (-1)^r\beta_1(\alpha_1\sigma_1 + \beta_1\tau_1))]\xi_r\phi_s \\ & + \gamma[\sigma_1(b_{21} - b_{12}) + \tau_1(b_{31} - b_{13})]\xi_r\psi_s, \end{aligned} \quad (4.63)$$

$$\begin{aligned}
G_2 = & (b_{32} - b_{23})(b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}) - (b_{21} - b_{12})(b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z}) \\
& + (\alpha_1\sigma_1 + \beta_1\tau_1)((-1)^s\sigma_1\gamma^2 + (-1)^r\alpha_1(\alpha_1\sigma_1 + \beta_1\tau_1))\xi_r\zeta_r\phi_s^2 \\
& + \gamma^2\tau_1(\alpha_1\tau_1 - \beta_1\sigma_1)\xi_r\zeta_r\psi_s^2 \\
& - (b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z})((-1)^s\sigma_1\gamma^2 + (-1)^r\alpha_1(\alpha_1\sigma_1 + \beta_1\tau_1))\xi_r\phi_s \\
& + (b_{32} - b_{23})\tau_1\gamma\xi_r\psi_s - (b_{21} - b_{12})(\alpha_1\sigma_1 + \beta_1\tau_1)\zeta_r\phi_s \\
& + (b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z})(\alpha_1\tau_1 - \beta_1\sigma_1)\gamma\zeta_r\psi_s,
\end{aligned} \tag{4.64}$$

$$\begin{aligned}
G_3 = & (b_{13} - b_{31})(b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z}) - (b_{32} - b_{23})(b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z}) \\
& + (\alpha_1\sigma_1 + \beta_1\tau_1)((-1)^s\tau_1\gamma^2 + (-1)^r\beta_1(\alpha_1\sigma_1 + \beta_1\tau_1))\xi_r\zeta_r\phi_s^2 \\
& - \gamma^2\sigma_1(\alpha_1\tau_1 - \beta_1\sigma_1)\xi_r\zeta_r\psi_s^2 \\
& - (b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z})((-1)^s\sigma_1\gamma^2 + (-1)^r\beta_1(\alpha_1\sigma_1 + \beta_1\tau_1))\xi_r\phi_s \\
& - (b_{32} - b_{23})\sigma_1\gamma\xi_r\psi_s - (b_{13} - b_{31})(\alpha_1\sigma_1 + \beta_1\tau_1)\zeta_r\phi_s \\
& - (b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z})(\alpha_1\tau_1 - \beta_1\sigma_1)\gamma\zeta_r\psi_s.
\end{aligned} \tag{4.65}$$

We assume that $\alpha_1\sigma_1 + \beta_1\tau_1 \neq 0$ and $\alpha_1\sigma + \beta_1\tau \neq 0$. Then the coefficients of $\zeta_r^2\phi_r^2$ in $\partial_{\mathcal{Y}}\Phi_1$ and $\partial_{\mathcal{X}}\Phi_2$ yield

$$\alpha_1\tau_1 = \beta_1\sigma_1. \tag{4.66}$$

For simplicity, we consider the special case in which

$$a_{21} = -a_{12} = \alpha', \quad a_{31} = -a_{13} = \beta' \sin \alpha, \quad a_{23} = a_{32} \quad \text{and} \quad B = B^t. \tag{4.67}$$

Furthermore, the nonlinear terms in $\Phi_{1\mathcal{Y}}$, $\Phi_{1\mathcal{Z}}$, $\Phi_{2\mathcal{X}}$, $\Phi_{2\mathcal{Z}}$, $\Phi_{3\mathcal{X}}$ and $\Phi_{3\mathcal{Y}}$ hint us that it is convenient to suppose

$$\gamma^2 = (-1)^{r+s+1}(\alpha_1^2 + \beta_1^2). \tag{4.68}$$

Thus the coefficients of $\xi_r\psi_s$ in $\Phi_{2\mathcal{Y}}$ and $\Phi_{2\mathcal{Z}}$ suggest that

$$\alpha_1\tau = \beta_1\sigma. \tag{4.69}$$

Hence we can write

$$\sigma_1 = g\sigma, \quad \tau_1 = g\tau. \tag{4.70}$$

As Xu did in [32], we get that

$$\begin{cases} \alpha_1 = \varphi\alpha', \\ \beta_1 = \varphi\beta' \sin \alpha \end{cases} \quad \text{and} \quad \begin{cases} \sigma = \mu\alpha', \\ \tau = \mu\beta' \sin \alpha \end{cases} \tag{4.71}$$

for some functions φ and μ . Moreover,

$$a_{22} = \frac{-a_{11}(\alpha')^2 - \alpha'\alpha'' + \beta' \sin \alpha (\beta' \sin \alpha)' + 2\alpha'\beta' \sin \alpha \beta' \cos \alpha}{(\alpha')^2 + (\beta' \sin \alpha)^2}, \tag{4.72}$$

$$a_{33} = \frac{-a_{11}(\beta' \sin \alpha)^2 + \alpha'\alpha'' - \beta' \sin \alpha (\beta' \sin \alpha)' - 2\alpha'\beta' \sin \alpha \beta' \cos \alpha}{(\alpha')^2 + (\beta' \sin \alpha)^2} \tag{4.73}$$

and

$$a_{23} = \frac{-a_{11}\alpha'\beta' \sin \alpha - \alpha''\beta' \sin \alpha - \alpha'\beta'' \sin \alpha - 2(\alpha')^2\beta' \cos \alpha + (\beta')^3 \sin^2 \alpha \cos \alpha}{(\alpha')^2 + (\beta' \sin \alpha)^2}. \tag{4.74}$$

The coefficients of $\zeta_r\psi_s$ and $\mathcal{Y}$ in $\Psi_1 = 0$ show that

$$b_{11} = b_{12} = b_{13} = 0. \tag{4.75}$$

Substituting (4.67)-(4.75) into (4.16), one gets that

$$a_{11} = -\frac{\gamma'}{\gamma} = \frac{(g(\alpha_1\sigma + \beta_1\tau))'}{g(\alpha_1\sigma + \beta_1\tau)}, \quad (4.76)$$

$$b_{22}(a_{23} + a_{32}) = b_{33}(a_{22} - a_{33}), \quad (4.77)$$

$$\begin{cases} b'_{22} = 2b_{23}\beta' \cos \alpha, \\ b'_{23} = -2b_{22}\beta' \cos \alpha, \end{cases} \quad (4.78)$$

$$\begin{cases} b_{22} = \frac{g}{\gamma^2}(\alpha_1\alpha'_1 - \beta_1\beta'_1 + \alpha_1^2 a_{22} - \beta_1\beta_1^2 a_{33} - 2\alpha_1\beta_1\beta' \cos \alpha), \\ b_{23} = \frac{g}{\gamma^2}((\alpha_1\beta_1)' + \alpha_1\beta_1(a_{22} + a_{33}) + (a_{23} + \beta' \cos \alpha)\tau^2\gamma g + (a_{23} - \beta' \cos \alpha)\sigma^2\gamma g) \end{cases} \quad (4.79)$$

and

$$\begin{cases} b_{22} = \frac{-\sigma(g\sigma\gamma)' + \tau(g\tau\gamma)' + a_{22}\sigma^2\gamma g - a_{33}\tau^2\gamma g + 2\sigma\tau\gamma g\beta' \cos \alpha}{\gamma(\sigma^2 + \tau^2)}, \\ b_{23} = \frac{-\tau(g\sigma\gamma)' + \sigma(g\tau\gamma)' - a_{11}\sigma\tau\gamma g + (a_{23} + \beta' \cos \alpha)\tau^2\gamma g + (a_{23} - \beta' \cos \alpha)\sigma^2\gamma g}{\gamma(\sigma^2 + \tau^2)}. \end{cases} \quad (4.80)$$

One shows that the above systems (4.77)-(4.80) are compatible with each other if and only if

$$\alpha''\beta' \sin \alpha - \alpha'\beta'' \sin \alpha - 2(\alpha')^2\beta' \cos \alpha - (\beta')^3 \sin^2 \alpha \cos \alpha = 0, \quad (4.81)$$

i.e.

$$\beta = \pm \int \frac{d\alpha}{\sin^2 \alpha \sqrt{d - \sin^{-2} \alpha}} + d_0 \quad (4.82)$$

for some constants d_0 and $d > 1$. Moreover, the system (4.79) says that

$$\begin{cases} b_{22} = \frac{g}{Q}(\frac{2\varphi'}{\varphi} + \frac{Q'}{2Q})((\alpha')^2 - (\beta' \sin \alpha)^2), \\ b_{23} = \frac{g}{Q}(\frac{4\varphi'}{\varphi} + \frac{Q'}{Q})\alpha'\beta' \sin \alpha, \end{cases} \quad (4.83)$$

where

$$Q = (\alpha')^2 + (\beta' \sin \alpha)^2. \quad (4.84)$$

Substituting (4.83) into (4.78), one finds that $g(\frac{2\varphi'}{\varphi} + \frac{Q'}{2Q})$ is a constant. We write

$$g(\frac{2\varphi'}{\varphi} + \frac{Q'}{2Q}) = 2c. \quad (4.85)$$

Thus

$$\varphi = k_1 Q^4 \exp\left(\int \frac{cdt}{g}\right) \quad (4.86)$$

with $k_1 \in \mathbb{R}$. Together with (4.76), one gets

$$\mu = k_2 Q^{-1} g^{-1} \exp\left(\int \frac{-2cdt}{g}\right) \quad (4.87)$$

with $k_2 \in \mathbb{R}$. Hence

$$\begin{aligned} a_{11} &= -\frac{Q'}{4Q} - \frac{c}{g} \\ &= -\frac{\alpha'\alpha'' + \beta' \sin \alpha (\beta' \sin \alpha)'}{2((\alpha')^2 + (\beta' \sin \alpha)^2)} - \frac{c}{g}. \end{aligned} \quad (4.88)$$

Theorem 4.4. Let α and g be arbitrary functions of t . The functions β , Q , φ , a_{22} , a_{33} , a_{23} , μ , ξ_r , ψ_r , ϕ_s and ψ_s are given by (4.82), (4.84), (4.72), (4.73), (4.74), (4.86), (4.87), (4.55), (4.56) and (4.57), respectively. Then we have the following solutions of the MHD equations (1.1)-(1.4):

$$\left\{ \begin{array}{l} u = - \left(\left(\frac{Q'}{4Q} - \frac{c}{g} \right) \cos \alpha + \alpha' \sin \alpha \right) \mathcal{X} - (\alpha' \cos \alpha + a_{22} \sin \alpha) \mathcal{Y} \\ \quad - (\beta' \cos \alpha + a_{23}) \mathcal{Z} \sin \alpha - \varphi \mu Q \zeta_r \phi_s \cos \alpha \mp \varphi \mu Q^{\frac{1}{2}} \xi_r \psi_s \alpha' \sin \alpha, \\ v = - \left(\frac{Q'}{4Q} - \frac{c}{g} \right) \sin \alpha \cos \beta + \alpha' \cos \alpha \cos \beta - \beta' \sin \alpha \sin \beta \mathcal{X} \\ \quad + (-\alpha' \sin \alpha \cos \beta + a_{22} \cos \alpha \cos \beta + a_{23} \sin \beta) \mathcal{Y} \\ \quad + (-\beta' \sin^2 \alpha \cos \beta + a_{23} \cos \alpha \cos \beta - a_{33} \sin \beta) \mathcal{Z} \\ \quad - \varphi \mu Q \zeta_r \phi_s \cos \beta \sin \alpha \pm (\alpha' \cos \alpha \cos \beta - \beta' \sin \alpha \sin \beta) \varphi \mu Q^{\frac{1}{2}} \xi_r \psi_s, \\ w = - \left(\frac{Q'}{4Q} - \frac{c}{g} \right) \sin \alpha \sin \beta + \alpha' \cos \alpha \sin \beta + \beta' \sin \alpha \cos \beta \mathcal{X} \\ \quad + (-\alpha' \sin \alpha \sin \beta + a_{22} \cos \alpha \sin \beta + a_{23} \cos \beta) \mathcal{Y} \\ \quad + (-\beta' \sin^2 \alpha \sin \beta + a_{23} \cos \alpha \sin \beta - a_{33} \cos \beta) \mathcal{Z} \\ \quad - \varphi \mu Q \zeta_r \phi_s \sin \alpha \sin \beta \pm (\alpha' \cos \alpha \sin \beta + \beta' \sin \alpha \cos \beta) \varphi \mu Q^{\frac{1}{2}} \xi_r \psi_s, \end{array} \right. \quad (4.89)$$

$$\left\{ \begin{array}{l} H^1 = - \frac{2c}{Q} \sin \alpha ((\alpha')^2 - (\beta' \sin \alpha)^2) \mathcal{Y} - \frac{4c}{Q} \alpha' \beta' \sin^2 \alpha \mathcal{Z} \\ \quad - g \varphi \mu Q \zeta_r \phi_s \cos \alpha \mp g \varphi \mu \alpha' Q^{\frac{1}{2}} \xi_r \psi_s \sin \alpha, \\ H^2 = \left(\frac{2c}{Q} \cos \alpha \cos \beta ((\alpha')^2 - (\beta' \sin \alpha)^2) - \frac{4c}{Q} \alpha' \beta' \sin \alpha \sin \beta \right) \mathcal{Y} \\ \quad + \left(\frac{4c}{Q} \alpha' \beta' \sin \alpha \cos \alpha \cos \beta + \frac{2c}{Q} \sin \beta ((\alpha')^2 - (\beta' \sin \alpha)^2) \right) \mathcal{Z} \\ \quad - g \varphi \mu Q \zeta_r \phi_s \sin \alpha \cos \beta \pm (\alpha' \cos \alpha \cos \beta - \beta' \sin \alpha \sin \beta) g \varphi \mu Q^{\frac{1}{2}} \xi_r \psi_s, \\ H^3 = \left(\frac{2c}{Q} \cos \alpha \sin \beta ((\alpha')^2 - (\beta' \sin \alpha)^2) + \frac{4c}{Q} \alpha' \beta' \sin \alpha \cos \beta \right) \mathcal{Y} \\ \quad + \left(\frac{4c}{Q} \alpha' \beta' \sin \alpha \cos \alpha \sin \beta - \frac{2c}{Q} \cos \beta ((\alpha')^2 - (\beta' \sin \alpha)^2) \right) \mathcal{Z} \\ \quad - g \varphi \mu Q \zeta_r \phi_s \sin \alpha \sin \beta \pm (\alpha' \cos \alpha \sin \beta + \beta' \sin \alpha \cos \beta) g \varphi \mu Q^{\frac{1}{2}} \xi_r \psi_s, \end{array} \right. \quad (4.90)$$

$$\begin{aligned} p = & -\rho((a'_{11} + a_{11}^2 - Q) \frac{\mathcal{X}^2}{2} + (a'_{22} + a_{22}^2 + a'_{23} - 2a_{23}\beta' \cos \alpha - (\alpha')^2) \frac{\mathcal{Y}^2}{2} \\ & (a'_{33} + a_{33}^2 + a'_{23} + 2a_{23}\beta' \cos \alpha - (\beta' \sin \alpha)^2) \frac{\mathcal{Z}^2}{2} \\ & - (\alpha' + 2\alpha'a_{22} + 2a_{23}\beta' \sin \alpha - (\beta')^2 \sin \alpha \cos \alpha) \mathcal{X} \mathcal{Y} \\ & - ((\beta' \sin \alpha)' + 2\alpha'a_{23} + \alpha'\beta' \cos \alpha) \mathcal{X} \mathcal{Z} \\ & + (a'_{23} - a_{11}a_{23} + (a_{22} - a_{33})\beta' \cos \alpha - \alpha'\beta' \sin \alpha) \mathcal{Y} \mathcal{Z} \\ & + (\delta_{r,1} + 4a\delta_{r,0}) \varphi^2 \mu^2 Q^2 \frac{\phi_r^2}{2} + (\delta_{s,1} + 4b\delta_{s,0}) \varphi^2 \mu^2 Q^2 \frac{\xi_r^2}{2} - 2\mu \xi_r \phi_s \\ & \pm (-1)^r \mu \sqrt{Q} \left(\frac{g'}{g} + \frac{2c}{g} + \frac{Q'}{2Q} \right) \zeta_r \psi_s \mp \varphi \mu \sqrt{Q} \left(\alpha' \left(\frac{\varphi'}{\varphi} + \frac{\alpha''}{\alpha'} + a_{22} \frac{\beta' \sin \alpha}{\alpha'} (a_{23} - \beta' \cos \alpha) \right) \mathcal{Y} \right. \\ & \left. + \beta' \sin \alpha \left(\frac{\varphi'}{\varphi} + \frac{(\beta' \sin \alpha)'}{\beta' \sin \alpha} + a_{33} + \frac{\alpha'}{\beta' \sin \alpha} (a_{23} + \beta' \cos \alpha) \mathcal{Z} \right) \xi_r \psi_s \right), \end{aligned} \quad (4.91)$$

where c is an arbitrary constant.

Set

$$\xi_r = \begin{cases} be^{\gamma_1 \mathcal{Y} + \gamma_2 \mathcal{Z}} - ce^{-(\gamma_1 \mathcal{Y} + \gamma_2 \mathcal{Z})} & \text{if } r = 1, \\ c \sin(\gamma_1 \mathcal{Y} + \gamma_2 \mathcal{Z} + b) & \text{if } r = 0 \end{cases} \quad (4.92)$$

and

$$\zeta_r = \begin{cases} be^{\gamma_1 \mathcal{Y} + \gamma_2 \mathcal{Z}} + ce^{-(\gamma_1 \mathcal{Y} + \gamma_2 \mathcal{Z})} & \text{if } r = 1, \\ c \cos(\gamma_1 \mathcal{Y} + \gamma_2 \mathcal{Z} + b) & \text{if } r = 0; \end{cases} \quad (4.93)$$

where γ_1 and γ_2 are functions of t , and b, c are real constants.

Suppose that

$$\begin{pmatrix} \mathcal{U} \\ \mathcal{V} \\ \mathcal{W} \end{pmatrix} = A \begin{pmatrix} \mathcal{X} \\ \mathcal{Y} \\ \mathcal{Z} \end{pmatrix} + \begin{pmatrix} f_1 - (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \zeta_r) \\ q_1 + \sigma \xi_r \\ s_1 + \tau \xi_r \end{pmatrix} \quad (4.94)$$

and

$$\begin{pmatrix} \mathcal{H}^1 \\ \mathcal{H}^2 \\ \mathcal{H}^3 \end{pmatrix} = B \begin{pmatrix} \mathcal{X} \\ \mathcal{Y} \\ \mathcal{Z} \end{pmatrix} + \begin{pmatrix} f_2 - (g_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \zeta_r) \\ q_2 + \sigma_1 \xi_r \\ s_2 + \tau_1 \xi_r \end{pmatrix}, \quad (4.95)$$

where $A = (a_{i,j})_{3 \times 3}$ and $B = (b_{i,j})_{3 \times 3}$ are 3×3 matrices whose entries are functions of t ; the functions $f_j, q_j, s_j, \sigma, \tau, \sigma_1$ and τ_1 are also functions of t to be determined.

The equations (4.16) become

$$\begin{aligned} & b'_{11} \mathcal{X} + b'_{12} \mathcal{Y} + b'_{13} \mathcal{Z} + f'_2 - (g'_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1)' \mathcal{X}) \zeta_r \\ & - (-1)^r (\gamma'_1 \mathcal{Y} + \gamma'_2 \mathcal{Z}) (g_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \mathcal{X}) \xi_r + (\alpha' \mathcal{Y} + \mathcal{Z} \beta' \sin \alpha) (b_{11} - (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \zeta_r) \\ & + ((a_{21} - \alpha') \mathcal{X} + a_{22} \mathcal{Y} + (a_{23} + \beta' \cos \alpha) \mathcal{Z} + q_1 + \sigma \xi_r) (b_{12} - (-1)^r \gamma_1 (g_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \mathcal{X}) \zeta_r) \\ & - ((\beta' \sin \alpha - a_{31}) \mathcal{X} + (\beta' \cos \alpha - a_{32}) \mathcal{Y} - a_{33} \mathcal{Z} - s_1 - \tau \xi_r) \\ & \times (b_{13} - (-1)^r \gamma_2 (g_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \mathcal{X}) \zeta_r) \\ & + ((a_{22} + a_{33}) + (\gamma_1 \sigma + \gamma_2 \tau) \zeta_r) (b_{11} \mathcal{X} + b_{12} \mathcal{Y} + b_{13} \mathcal{Z} + f_2 - (g_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \mathcal{X}) \zeta_r) \\ & - (\alpha' + a_{12} - (-1)^r \gamma_1 (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) \xi_r) (b_{21} \mathcal{X} + b_{22} \mathcal{Y} + b_{23} \mathcal{Z} + q_2 + \sigma_1 \xi_r) \\ & - (\beta' \sin \alpha + a_{13} - (-1)^r \gamma_2 (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) \xi_r) (b_{31} \mathcal{X} + b_{32} \mathcal{Y} + b_{33} \mathcal{Z} + s_2 + \sigma_2 \xi_r) \\ & - (-b_{11} + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \zeta_r) (a_{11} \mathcal{X} + a_{12} \mathcal{Y} + a_{13} \mathcal{Z} + f_1 - (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) \xi_r) \\ & + (-1)^r \eta (\gamma_1^2 + \gamma_2^2) (g_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \mathcal{X}) \zeta_r = 0, \end{aligned} \quad (4.96)$$

$$\begin{aligned} & b'_{21} \mathcal{X} + b'_{22} \mathcal{Y} + b'_{23} \mathcal{Z} + q'_2 + \sigma'_1 \xi_r + (\gamma'_1 \mathcal{Y} + \gamma'_2 \mathcal{Z}) \sigma_1 \xi_r \\ & + b_{21} (a_{11} \mathcal{X} + (a_{12} + \alpha') \mathcal{Y} + (a_{13} + \beta' \sin \alpha) \mathcal{Z} + f_1 - (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) \xi_r) \\ & + (b_{23} + \gamma_2 \sigma_1 \xi_r) ((a_{31} - \beta' \sin \alpha) \mathcal{X} + (a_{32} - \beta' \cos \alpha) \mathcal{Y} + a_{33} \mathcal{Z} + s_1 + \tau \xi_r) \\ & + (\alpha' - a_{21}) (b_{11} \mathcal{X} + b_{12} \mathcal{Y} + b_{13} \mathcal{Z} + f_2 - (g_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \mathcal{X}) \zeta_r) \\ & - (a_{22} + \gamma_1 \sigma \xi_r) (b_{21} \mathcal{X} + b_{22} \mathcal{Y} + b_{23} \mathcal{Z} + q_2 + \sigma_1 \xi_r) + (b_{22} + \gamma_1 \sigma_1 \xi_r) (-\alpha' \mathcal{X} + \mathcal{Z} \beta' \cos \alpha) \\ & - (a_{23} + \beta' \cos \alpha + \gamma_2 \sigma \xi_r) (b_{31} \mathcal{X} + b_{32} \mathcal{Y} + b_{33} \mathcal{Z} + s_2 + \tau_1 \xi_r) \\ & + (b_{22} + \gamma_1 \sigma_1 \xi_r) (a_{21} \mathcal{X} + a_{22} \mathcal{Y} + a_{23} \mathcal{Z} + q_1 + \sigma \xi_r) - (-1)^r \eta (\gamma_1^2 + \gamma_2^2) \sigma_1 \xi_r = 0 \end{aligned} \quad (4.97)$$

and

$$\begin{aligned} & b'_{31} \mathcal{X} + b'_{32} \mathcal{Y} + b'_{33} \mathcal{Z} + s'_2 + \tau'_1 \xi_r + (\gamma'_1 \mathcal{Y} + \gamma'_2 \mathcal{Z}) \tau_1 \xi_r \\ & + b_{31} (a_{11} \mathcal{X} + (a_{12} + \alpha') \mathcal{Y} + (a_{13} + \beta' \sin \alpha) \mathcal{Z} + f_1 - (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) \xi_r) \\ & + (b_{32} + \gamma_1 \tau_1 \xi_r) ((a_{21} - \alpha') \mathcal{X} + a_{22} \mathcal{Y} + (a_{23} + \beta' \cos \alpha) \mathcal{Z} + q_1 + \sigma \xi_r) \\ & + (b_{33} + \gamma_2 \tau_1 \xi_r) ((a_{31} - \beta' \sin \alpha) \mathcal{X} + (a_{32} - \beta' \cos \alpha) \mathcal{Y} + a_{33} \mathcal{Z} + s_1 + \tau \xi_r) \\ & + (\beta' \sin \alpha - a_{31}) (b_{11} \mathcal{X} + b_{12} \mathcal{Y} + b_{13} \mathcal{Z} + f_2 - (g_2 + (\gamma_1 \sigma_1 + \gamma_2 \tau_1) \mathcal{X}) \zeta_r) \\ & + (\beta' \cos \alpha - a_{32} - \gamma_1 \tau \xi_r) (b_{21} \mathcal{X} + b_{22} \mathcal{Y} + b_{23} \mathcal{Z} + q_2 + \sigma_1 \xi_r) \\ & - (a_{33} + \gamma_2 \tau \xi_r) (b_{31} \mathcal{X} + b_{32} \mathcal{Y} + b_{33} \mathcal{Z} + s_2 + \tau_1 \xi_r) - (-1)^r \eta (\gamma_1^2 + \gamma_2^2) \tau_1 \xi_r = 0. \end{aligned} \quad (4.98)$$

The coefficients of $\xi_r \zeta_r$ and ξ_r^2 say that

$$\begin{cases} \tau_1 \sigma - \tau \sigma_1 = 0, \\ (\gamma_1 \sigma_1 + \gamma_2 \tau_1) g_1 = (\gamma_1 \sigma + \gamma_2 \tau) g_2. \end{cases} \quad (4.99)$$

Moreover, the coefficients of $\xi_r \zeta_r$ in $\Phi_{1\mathcal{Z}} = \Phi_{3\mathcal{X}}$ yield

$$\gamma_1 \tau = \gamma_2 \sigma. \quad (4.100)$$

The above two equations suggest that

$$\begin{cases} \sigma_1 = h\sigma, \\ \tau_1 = h\tau, \end{cases} \quad (4.101)$$

where

$$h = \frac{g_2}{g_1}. \quad (4.102)$$

Motivated by Theorem 4.4, we assume that

$$a_{21} = -a_{12} = \alpha', \quad a_{31} = -a_{13} = \beta' \sin \alpha. \quad (4.103)$$

The coefficients of ξ_r in $\Phi_{1\mathcal{Y}} = \Phi_{2\mathcal{X}}$ and the coefficients of $\mathcal{Y}$ in (4.96) show that

$$b_{11} = b_{12} = b_{13} = 0. \quad (4.104)$$

The coefficients of ξ_r and ζ_r in (4.96)-(4.98) yield

$$b_{21} = b_{31} = 0. \quad (4.105)$$

Now

$$\begin{aligned} R_1 = & (a'_{11} + a_{11}^2 + (\beta' \sin \alpha)^2) \mathcal{X} - (\alpha'' + 2\alpha' a_{22} - 2a_{32} \beta' \sin \alpha + (\beta')^2 \sin \alpha \cos \alpha) \mathcal{Y} \\ & + (-(\beta' \sin \alpha)' - \alpha' (a_{23} + \beta' \cos \alpha) - 2a_{33} \beta' \sin \alpha - \alpha' a_{23}) \mathcal{Z} \\ & + \{-2\alpha' \sigma - 2\beta' \tau \sin \alpha - (-1)^r (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) [(\gamma'_1 + \gamma_1 a_{22} + \gamma_2 (a_{32} - \beta' \cos \alpha)) \mathcal{Y} \\ & + (\gamma'_2 + \gamma_1 (a_{23} + \beta' \cos \alpha) - 2a_{33} \beta' \sin \alpha) \mathcal{Z} + \gamma_1 q_1 + \gamma_2 s_1]\} \xi_r + \{-(g'_1 + (\gamma_1 \sigma + \gamma_2 \tau)' \mathcal{X}) \\ & - a_{11} (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) - (a_{11} \mathcal{X} + f_1) (\gamma_1 \sigma + \gamma_2 \tau) \\ & + (-1)^r \nu (\gamma_1^2 + \gamma_2^2) (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X})\} \zeta_r \\ & + (4bc\delta_{r,0} + c^2 \delta_{r,1}) (\gamma_1 \sigma + \gamma_2 \tau) (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) \\ & + f'_1 + a_{11} f_1 - 2\alpha' q_1 - 2s_1 \beta' \sin \alpha, \end{aligned} \quad (4.106)$$

$$\begin{aligned} R_2 = & (\alpha'' + 2\alpha' a_{11} - (\beta')^2 \sin \alpha \cos \alpha) \mathcal{X} \\ & + (a'_{22} + a_{22}^2 + a_{23} (a_{32} - \beta' \cos \alpha) - (\alpha')^2 - a_{32} \beta' \cos \alpha) \mathcal{Y} \\ & + (a'_{23} + a_{22} (a_{23} + \beta' \cos \alpha) + a_{23} a_{33} - \alpha' \beta' \sin \alpha - a_{33} \beta' \cos \alpha) \mathcal{Z} \\ & + (\sigma' + a_{22} \sigma + (a_{23} - \beta' \cos \alpha) \tau - (-1)^r \nu (\gamma_1^2 + \gamma_2^2) \sigma) \xi_r \\ & + \{-2\alpha' (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) + \sigma [(\gamma'_1 + \gamma_1 a_{22} + \gamma_2 (a_{32} - \beta' \cos \alpha)) \mathcal{Y} \\ & + (\gamma'_2 + \gamma_1 (a_{23} + \beta' \cos \alpha) - 2a_{33} \beta' \sin \alpha) \mathcal{Z} + \gamma_1 q_1 + \gamma_2 s_1]\} \zeta_r \\ & + \sigma (\gamma_1 \sigma + \gamma_2 \tau) \xi_r \zeta_r + q'_1 + a_{22} q_1 + (a_{23} - \beta' \cos \alpha) s_1, \end{aligned} \quad (4.107)$$

$$\begin{aligned} R_3 = & ((\beta' \sin \alpha)' + 2a_{11} \beta' \sin \alpha + \alpha' \beta' \cos \alpha) \mathcal{X} \\ & + (a'_{32} + a_{32} a_{22} + a_{33} (a_{32} - \beta' \cos \alpha) - \alpha' \beta' \sin \alpha - a_{22} \beta' \cos \alpha) \mathcal{Y} \\ & + (a'_{33} + a_{32} (a_{23} + \beta' \cos \alpha) + a_{33}^2 - (\beta' \sin \alpha)^2 + a_{23} \beta' \cos \alpha) \mathcal{Z} \\ & + (\tau' + a_{33} \tau + (a_{32} + \beta' \cos \alpha) \sigma - (-1)^r \nu (\gamma_1^2 + \gamma_2^2) \tau) \xi_r \\ & + \{-2(g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) \beta' \sin \alpha + \tau [(\gamma'_1 + \gamma_1 a_{22} + \gamma_2 (a_{32} - \beta' \cos \alpha)) \mathcal{Y} \\ & + (\gamma'_2 + \gamma_1 (a_{23} + \beta' \cos \alpha) - 2a_{33} \beta' \sin \alpha) \mathcal{Z} + \gamma_1 q_1 + \gamma_2 s_1]\} \zeta_r \\ & + \tau (\gamma_1 \sigma + \gamma_2 \tau) \xi_r \zeta_r + s'_1 + a_{33} s_1 + (a_{32} + \beta' \cos \alpha) q_1 + 2f_1 \beta' \sin \alpha, \end{aligned} \quad (4.108)$$

$$G_1 = (-1)^r h^2 (\gamma_1 \sigma + \gamma_2 \tau) (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) \xi_r^2 + (-1)^r h (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X}) (\gamma_1 b_{23} \mathcal{Z} + \gamma_2 b_{32} \mathcal{Y} + \gamma_1 q_2 + \gamma_2 s_2) \xi_r, \quad (4.109)$$

$$G_2 = b_{32} (b_{32} - b_{23}) \mathcal{Y} + h^2 (-1)^r \gamma_1 (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X})^2 \xi_r \zeta_r + h ((b_{32} - b_{23}) \tau - (-1)^r f_2 \gamma_1 (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X})) \xi_r + (b_{32} - b_{23}) s_2, \quad (4.110)$$

$$G_3 = b_{23} (b_{32} - b_{23}) \mathcal{Z} + h^2 (-1)^r \gamma_2 (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X})^2 \xi_r \zeta_r - h ((b_{32} - b_{23}) \sigma - (-1)^r f_2 \gamma_1 (g_1 + (\gamma_1 \sigma + \gamma_2 \tau) \mathcal{X})) \xi_r. \quad (4.111)$$

By (4.18), we have that

$$\begin{cases} a_{33} = \frac{\alpha''}{\alpha} - \frac{(\beta')^2 \sin \alpha \cos \alpha}{\alpha'} + \frac{\beta' \sin \alpha a_{32}}{\alpha'}, \\ a_{22} = \frac{(\beta' \sin \alpha)'}{\beta' \sin \alpha} + \frac{\alpha' \cos \alpha}{\sin \alpha} + \frac{\alpha' a_{23}}{\beta' \sin \alpha} \end{cases}, \quad (4.112)$$

and

$$(a_{23} - a_{32}) a_{11} = a'_{23} - a'_{32}. \quad (4.113)$$

The coefficients of ζ_r in $\Phi_{1y} = \Phi_{2x}$ say that

$$\begin{cases} b_{32} = -\frac{\gamma'_1 + \gamma_1 a_{22} + \gamma_2 (a_{32} - \beta' \cos \alpha)}{\mu_0 h \gamma_2}, \\ b_{23} = -\frac{\gamma'_2 + \gamma_1 (a_{23} + \beta' \cos \alpha) + \gamma_2 a_{33}}{\mu_0 h \gamma_1}. \end{cases} \quad (4.114)$$

But the coefficients of ξ in (4.96) yield

$$\begin{cases} b_{32} = \frac{h(\gamma'_1 + \gamma_1 a_{22} + \gamma_2 (a_{32} - \beta' \cos \alpha))}{\gamma_2}, \\ b_{23} = \frac{h(\gamma'_2 + \gamma_1 (a_{23} + \beta' \cos \alpha) + \gamma_2 a_{33})}{\gamma_1}. \end{cases} \quad (4.115)$$

Thus

$$b_{23} = b_{32} = 0 \quad (4.116)$$

and

$$\begin{cases} \gamma'_1 + \gamma_1 a_{22} + \gamma_2 (a_{32} - \beta' \cos \alpha) = 0, \\ \gamma'_2 + \gamma_1 (a_{23} + \beta' \cos \alpha) + \gamma_2 a_{33} = 0. \end{cases} \quad (4.117)$$

Again by (4.18), we have that

$$\begin{cases} \frac{(\gamma_1 \sigma + \gamma_2 \tau)'}{\gamma_1 \sigma + \gamma_2 \tau} + 2a_{11} - (-1)^r \nu (\gamma_1^2 + \gamma_2^2) = 0, \\ \frac{g'_1}{g_1} + a_{11} + \frac{f_1 (\gamma_1 \sigma + \gamma_2 \tau)}{g_1} - (-1)^r \nu (\gamma_1^2 + \gamma_2^2) + \frac{(-1)^r \mu_0 f_2 h (\gamma_1 \sigma + \gamma_2 \tau)}{g_1} = 0, \end{cases} \quad (4.118)$$

$$\gamma_1 \beta' \sin \alpha = \alpha' \gamma_2 \quad (4.119)$$

and

$$\gamma_1 q_1 + \gamma_2 s_1 + \mu_0 h (\gamma_1 q_2 + \gamma_2 s_2) = 0. \quad (4.120)$$

Since

$$h(\gamma_1 q_1 + \gamma_2 s_1) + \gamma_1 q_2 + \gamma_2 s_2 = 0, \quad (4.121)$$

comparing with (4.96), we get

$$\gamma_1 q_1 + \gamma_2 s_1 = \gamma_1 q_2 + \gamma_2 s_2 = 0. \quad (4.122)$$

Furthermore, by (4.100) and (4.119), we can write

$$\begin{cases} \gamma_1 = \varphi\alpha' \\ \gamma_2 = \varphi\beta' \sin \alpha \end{cases} \quad \text{and} \quad \begin{cases} \sigma = \mu\alpha' \\ \tau = \mu\beta' \sin \alpha \end{cases} \quad (4.123)$$

for some functions φ and μ .

From (4.96)-(4.98), we have that

$$\begin{cases} (h(\gamma_1\sigma + \gamma_2\tau))' - (-1)^r \eta h(\gamma_1^2 + \gamma_2^2)g_1 = 0 \\ (hg_1)' - ha_{11}g_1 + hf_1(\gamma_1\sigma + \gamma_2\tau) - f_2(\gamma_1\sigma + \gamma_2\tau) - \eta(-1)^r \eta h(\gamma_1^2 + \gamma_2^2)g_1 = 0, \end{cases} \quad (4.124)$$

$$\begin{cases} (\sigma h)' - a_{12}h\sigma - h\tau(a_{23} + \beta' \cos \alpha) - (-1)^r \eta h(\gamma_1^2 + \gamma_2^2)\sigma = 0 \\ (\tau h)' + (\beta' \cos \alpha - a_{32})h\sigma - a_{33}h\tau - (-1)^r \eta h(\gamma_1^2 + \gamma_2^2)\tau = 0 \end{cases} \quad (4.125)$$

and

$$\begin{cases} f_2' - a_{11}f_2 = 0 \\ q_2' - a_{22}q_2 - (a_{23} + \beta' \cos \alpha)s_2 = 0 \\ s_2' + (\beta' \cos \alpha - a_{32})q_2 - a_{33}s_2 = 0. \end{cases} \quad (4.126)$$

For simplicity, we assume that

$$f_2 = q_2 = s_2 = 0. \quad (4.127)$$

Then by (4.112) and (4.117), one has

$$\begin{cases} a_{22} = \frac{(\beta' \sin \alpha)'}{\beta' \sin \alpha} + \frac{\alpha' \beta' \cos \alpha}{\beta' \sin \alpha} + \frac{\alpha' a_{23}}{\beta' \sin \alpha} \\ a_{33} = -\frac{\varphi'}{\varphi} - \frac{(\beta' \sin \alpha)'}{\beta' \sin \alpha} - \frac{\alpha' \beta' \cos \alpha}{\beta' \sin \alpha} - \frac{\alpha' a_{23}}{\beta' \sin \alpha} \\ a_{32} = -\frac{\alpha'}{\beta' \sin \alpha} \left(\frac{\varphi'}{\varphi} + \frac{\alpha''}{\alpha} + \frac{(\beta' \sin \alpha)'}{\beta' \sin \alpha} + \frac{\alpha' \beta' \cos \alpha}{\beta' \sin \alpha} \right) + \beta' \cos \alpha - \frac{(\alpha')^2 a_{23}}{(\beta' \sin \alpha)^2}. \end{cases} \quad (4.128)$$

Thus

$$a_{11} = \frac{\varphi'}{\varphi}. \quad (4.129)$$

The system (4.118) says that

$$3\frac{\varphi'}{\varphi} + \frac{\mu'}{\mu} + \frac{Q'}{Q} = (-1)^r \nu \varphi Q, \quad (4.130)$$

where

$$Q = (\alpha')^2 + (\beta' \sin \alpha)^2. \quad (4.131)$$

Then we get

$$\mu = \frac{c_1}{\varphi^3 Q} \exp\left(\int (-1)^r \nu \varphi Q dt\right) \quad (4.132)$$

for some constants c_1 .

We consider the special case of (4.128), in which

$$a_{23} = a_{32}. \quad (4.133)$$

Hence we have

$$a_{23} = -\frac{\alpha' \beta' \sin \alpha}{Q} \left(\frac{\varphi'}{\varphi} + \frac{\alpha''}{\alpha} + \frac{(\beta' \sin \alpha)'}{\beta' \sin \alpha} + \frac{\alpha' \beta' \cos \alpha}{\beta' \sin \alpha} - \frac{(\beta')^2 \sin \alpha \cos \alpha}{\alpha'} \right). \quad (4.134)$$

One shows that the system (4.124) is compatible by (4.118). Moreover, the equation (4.125)₁ is compatible with (4.124) if and only if

$$\alpha'' \beta' \sin \alpha - \alpha' \beta'' \sin \alpha - 2(\alpha')^2 \beta' \cos \alpha - (\beta')^3 \sin^2 \alpha \cos \alpha = 0, \quad (4.135)$$

i.e.

$$\beta = \pm \int \frac{d\alpha}{\sin^2 \alpha \sqrt{d - \sin^{-2} \alpha}} + d_0 \quad (4.136)$$

for some constants d_0 and $d > 1$. In this case, one shows the system (4.125) is compatible. Thus one gets that

$$h = c_2 \varphi^2 \exp((-1)^r \int (\eta - \nu) \varphi Q dt), \quad (4.137)$$

where c_2 is a real constant.

Write

$$\begin{cases} s_1 = \lambda \alpha', \\ q_1 = \lambda \beta' \sin \alpha. \end{cases} \quad (4.138)$$

Then we have

$$\begin{aligned} p &= -\rho(-(-1)^r (c_2 \varphi^2 \exp((-1)^r \int (\eta - \nu) \varphi Q dt))^2 \mu_0 ((g_1 + \varphi \mu Q \mathcal{X})^2 \xi_r^2) \\ &+ \left(\frac{\varphi''}{\varphi} - Q \right) \frac{\mathcal{X}^2}{2} + (a'_{22} + a_{22}^2 + a_{23}^2 - 2a_{23} \beta' \cos \alpha - (\alpha')^2) \frac{\mathcal{Y}^2}{2} \\ &+ (a'_{33} + a'_{23} + a_{33}^2 + 2a_{23} \beta' \cos \alpha - (\beta' \sin \alpha)^2) \frac{\mathcal{Z}^2}{2} + (\alpha'' + 2\alpha' \frac{\varphi'}{\varphi} - (\beta')^2 \sin \alpha \cos \alpha) \mathcal{X} \mathcal{Y} \\ &+ ((\beta' \sin \alpha)' + 2\frac{\varphi'}{\varphi} \beta' \sin \alpha + \alpha' \beta' \cos \alpha) \mathcal{X} \mathcal{Z} \\ &+ (a_{23} - a_{23} \frac{\varphi'}{\varphi} + (a_{22} - a_{33}) \beta' \cos \alpha - \alpha' \beta' \cos \alpha) \mathcal{Y} \mathcal{Z} \\ &+ \frac{1}{2} (4bc\delta_{r,0} + c^2 \delta_{r,1}) (g_1 + \varphi \mu Q \mathcal{X})^2 + (f'_1 + \frac{\varphi' f_1}{f_1} - 4\lambda \alpha' \beta' \sin \alpha) \mathcal{X} \\ &+ ((\lambda \beta \sin \alpha)' + a_{22} \lambda \beta \sin \alpha + (a_{23} - \beta' \cos \alpha) (\lambda \alpha')) \mathcal{Y} \\ &+ ((\lambda \alpha')' + 2f_1 \beta' \sin \alpha + (a_{23} + \beta' \cos \alpha) \lambda \beta' \sin \alpha + a_{33} \lambda \alpha') \mathcal{Z} \\ &- 2\mu Q \mathcal{X} \xi_r - 2\frac{\xi_r}{\varphi} + (-1)^r \frac{\mu}{\varphi} \left(\frac{\mu'}{\mu} - \frac{\varphi'}{\varphi} - (-1)^r \nu \varphi^2 Q \right) \zeta_r \end{aligned} \quad (4.139)$$

modulo the transformation in (1.9), where c_2 is an arbitrary constant.

Theorem 4.5. *Let $\alpha, \varphi, g_1, \lambda, f_1$ be functions of t . The functions $\mu, \xi_r, \zeta_r, \beta, Q, a_{22}, a_{33}, a_{23}$ are given by (4.132), (4.92), (4.93), (4.136), (4.131), (4.128) and (4.134), respectively. Then we have the following solutions of the MHD equations (1.1)-(1.4):*

$$\left\{ \begin{aligned} u &= \left(\frac{\varphi'}{\varphi} \cos \alpha - \alpha' \sin \alpha \right) \mathcal{X} - (\alpha' \cos \alpha + a_{22} \sin \alpha) \mathcal{Y} - (\beta' \cos \alpha - a_{23}) \mathcal{Z} \sin \alpha \\ &\quad + f_1 \cos \alpha - \lambda \beta' \sin^2 \alpha - (g_1 + \varphi \mu Q \mathcal{X}) \zeta_r \cos \alpha - \mu \xi_r \alpha' \sin \alpha, \\ v &= \left(\frac{\varphi'}{\varphi} \sin \alpha \cos \beta + \alpha' \cos \alpha \cos \beta - \beta' \sin \alpha \sin \beta \right) \mathcal{X} \\ &\quad + (-\alpha' \sin \alpha \cos \beta + a_{22} \cos \alpha \cos \beta - a_{23} \sin \beta) \mathcal{Y} \\ &\quad + (-\beta' \sin^2 \alpha \cos \beta + a_{23} \cos \alpha \cos \beta - a_{33} \sin \beta) \mathcal{Z} \\ &\quad + f_1 \sin \alpha \cos \beta + \lambda \beta' \sin \alpha \cos \alpha \cos \beta - \lambda \alpha' \sin \beta \\ &\quad - (g_1 + \varphi \mu Q \mathcal{X}) \zeta_r \sin \alpha \cos \beta + \mu \alpha' \xi_r \cos \alpha \cos \beta - \mu \beta' \xi_r \sin \alpha \sin \beta, \\ w &= \left(\frac{\varphi'}{\varphi} \sin \alpha \sin \beta + \alpha' \cos \alpha \sin \beta + \beta' \sin \alpha \cos \beta \right) \mathcal{X} \\ &\quad + (-\alpha' \sin \alpha \sin \beta + a_{22} \cos \alpha \sin \beta + a_{23} \cos \beta) \mathcal{Y} \\ &\quad + (-\beta' \sin^2 \alpha \sin \beta + a_{23} \cos \alpha \sin \beta + a_{33} \cos \beta) \mathcal{Z} \\ &\quad + f_1 \sin \alpha \sin \beta + \lambda \beta' \sin \alpha \cos \alpha \sin \beta + \lambda \alpha' \cos \beta \\ &\quad - (g_1 + \varphi \mu Q \mathcal{X}) \zeta_r \sin \alpha \sin \beta + \mu \alpha' \xi_r \cos \alpha \sin \beta + \mu \beta' \xi_r \sin \alpha \cos \beta, \end{aligned} \right. \quad (4.140)$$

$$\left\{ \begin{array}{l} H^1 = -c_2 \varphi^2 \exp((-1)^r \int (\eta - \nu) \varphi Q dt) ((g_1 + \varphi \mu Q \mathcal{X}) \zeta_r \cos \alpha + \mu \alpha' \xi_r \sin \alpha), \\ H^2 = c_2 \varphi^2 \exp((-1)^r \int (\eta - \nu) \varphi Q dt) (-(g_1 + \varphi \mu Q \mathcal{X}) \zeta_r \sin \alpha \cos \beta \\ \quad + \mu \alpha' \xi_r \cos \alpha \cos \beta - \mu \beta' \xi_r \sin \alpha \sin \beta), \\ H^3 = c_2 \varphi^2 \exp((-1)^r \int (\eta - \nu) \varphi Q dt) (-(g_1 + \varphi \mu Q \mathcal{X}) \zeta_r \sin \alpha \sin \beta \\ \quad + \mu \alpha' \xi_r \cos \alpha \sin \beta + \mu \beta' \xi_r \sin \alpha \cos \beta) \end{array} \right. \quad (4.141)$$

and p is given by (4.139).

Set

$$\phi_r = \begin{cases} e^{a_1 \mathcal{Y}} - a e^{-a_1 \mathcal{Y}} & \text{if } r = 0, \\ \sin(a_1 \mathcal{Y}) & \text{if } r = 1, \end{cases} \quad \psi_r = \begin{cases} e^{a_1 \mathcal{Y}} + a e^{-a_1 \mathcal{Y}} & \text{if } r = 0, \\ \cos(a_1 \mathcal{Y}) & \text{if } r = 1, \end{cases} \quad (4.142)$$

$$\xi_r = \begin{cases} b e^{a_1 \mathcal{Y}} - c e^{-a_1 \mathcal{Y}} & \text{if } r = 0, \\ c \sin(a_1 \mathcal{Y} + b) & \text{if } r = 1, \end{cases} \quad \zeta_r = \begin{cases} b e^{a_1 \mathcal{Y}} + c e^{-a_1 \mathcal{Y}} & \text{if } r = 0, \\ c \cos(a_1 \mathcal{Y} + b) & \text{if } r = 1 \end{cases} \quad (4.143)$$

and

$$\theta_r = \begin{cases} b_1 e^{a_1 \mathcal{Y}} - c_1 e^{-a_1 \mathcal{Y}} & \text{if } r = 0, \\ c_1 \sin(a_1 \mathcal{Y} + b_1) & \text{if } r = 1, \end{cases} \quad \varepsilon_r = \begin{cases} b_1 e^{a_1 \mathcal{Y}} + c_1 e^{-a_1 \mathcal{Y}} & \text{if } r = 0, \\ c_1 \cos(a_1 \mathcal{Y} + b_1) & \text{if } r = 1, \end{cases} \quad (4.144)$$

where a_1 is a functions of t and a, b, b_1, c, c_1 are real constants .

We assume that

$$\left\{ \begin{array}{l} \mathcal{U} = a_{11} \mathcal{X} + a_{12} \mathcal{Y} + a_{13} \mathcal{Z} + a_1 f(\theta_r + a_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 a_2 \mathcal{X}^2 \psi_r), \\ \mathcal{V} = a_{21} \mathcal{X} + a_{22} \mathcal{Y} + a_{23} \mathcal{Z} + f(\xi_r + 2a_1 a_2 \mathcal{X} \phi_r), \\ \mathcal{W} = a_{31} \mathcal{X} + a_{32} \mathcal{Y} + a_{33} \mathcal{Z} \end{array} \right. \quad (4.145)$$

and

$$\left\{ \begin{array}{l} \mathcal{H}^1 = b_{11} \mathcal{X} + b_{12} \mathcal{Y} + b_{13} \mathcal{Z} + a_1 g(\theta_r + b_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 b_2 \mathcal{X}^2 \psi_r), \\ \mathcal{V} = b_{21} \mathcal{X} + b_{22} \mathcal{Y} + b_{23} \mathcal{Z} + g(\xi_r + 2a_1 b_2 \mathcal{X} \phi_r), \\ \mathcal{W} = b_{31} \mathcal{X} + b_{32} \mathcal{Y} + b_{33} \mathcal{Z}, \end{array} \right. \quad (4.146)$$

where $a_{ij}, b_{ij}, a_2 (\neq 0), b_2 (\neq 0), f (\neq 0)$ and $g (\neq 0)$ are functions of t , and

$$\left\{ \begin{array}{l} a_{11} + a_{22} + a_{33} = 0, \\ b_{11} + b_{22} + b_{33} = 0. \end{array} \right. \quad (4.147)$$

Substituting (4.145)-(4.147) into (4.16) and (4.18), we get that

$$\begin{aligned} R_1 &= a'_{11} \mathcal{X} + a'_{12} \mathcal{Y} + a'_{13} \mathcal{Z} + (a_1 f)' (\theta_r + a_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 a_2 \mathcal{X}^2 \psi_r) \\ &+ a_1 f (a'_1 \mathcal{Y} \varepsilon_r + a'_2 \mathcal{Y} \phi_r + a'_1 a_2 \mathcal{Y}^2 \psi_r - (-1)^r a'_1 \mathcal{X} \mathcal{Y} \xi_r - (a_1 a_2)' \mathcal{X} \psi_r - (-1)^r a'_1 a_1 a_2 \mathcal{X}^2 \mathcal{Y} \phi_r) \\ &+ (a_{11} - a_1 f(\zeta_r + 2a_1 a_2 \mathcal{X} \psi_r)) (a_{11} \mathcal{X} + (a_1 2 + \alpha') \mathcal{Y} + (a_{13} + \beta' \sin \alpha) \mathcal{Z} \\ &+ a_1 f(\theta_r + a_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 a_2 \mathcal{X}^2 \psi_r)) \\ &+ (a_{12} + a_1 f(a_1 \varepsilon_r + a_2 \phi_r + a_1 a_2 \mathcal{Y} \psi_r - (-1)^r a_1 \mathcal{X} \xi_r - (-1)^r a_1^2 a_2 \mathcal{X}^2 \phi_r)) \\ &\times ((a_{21} - \alpha') \mathcal{X} + a_{22} \mathcal{Y} + (a_{23} + \beta' \cos \alpha) \mathcal{Z} + f(\xi_r + 2a_1 a_2 \mathcal{X} \phi_r)) \\ &+ a_{13} ((a_{31} - \beta' \sin \alpha) \mathcal{X} + (a_{32} - \beta' \cos \alpha) \mathcal{Y} + a_{33} \mathcal{Z}) \\ &- \alpha' (a_{21} \mathcal{X} + a_{22} \mathcal{Y} + a_{23} \mathcal{Z} + f(\xi_r + 2a_1 a_2 \mathcal{X} \phi_r)) - (a_{31} \mathcal{X} + a_{32} \mathcal{Y} + a_{33} \mathcal{Z}) \beta' \sin \alpha \\ &- (-1)^r \nu a_1^3 f(\theta_r + a_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 a_2 \mathcal{X}^2 \psi_r), \end{aligned} \quad (4.148)$$

$$\begin{aligned}
G_1 &= (b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z} + g(\xi_r + 2a_1b_2\mathcal{X}\phi_r)) \\
&\times (b_{21} - b_{12} + a_1g(-a_1\varepsilon_r + b_2\phi_r - a_1b_2\mathcal{Y}\psi_r + (-1)^r a_1(\mathcal{X}\xi_r + a_1b_2\mathcal{X}^2\phi_r)) \\
&- (b_{13} - b_{31})(b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}), \tag{4.149}
\end{aligned}$$

$$\begin{aligned}
R_2 &= a'_{21}\mathcal{X} + a'_{22}\mathcal{Y} + a'_{23}\mathcal{Z} + f'(\xi_r + 2a_1a_2\mathcal{X}\phi_r) \\
&+ f(a'_1\mathcal{Y}\zeta_r + 2(a_1a_2)'\mathcal{X}\phi_r + 2a'_1a_1a_2\mathcal{X}\mathcal{Y}\psi_r) \\
&+ (a_{21} + 2a_1a_2f\phi_r)(a_{11}\mathcal{X} + (a_{12} + \alpha')\mathcal{Y} + (a_{13} + \beta'\sin\alpha)\mathcal{Z} \\
&+ a_1f(\theta_r + a_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1a_2\mathcal{X}^2\psi_r)) \\
&+ (a_{22} + a_1f(\zeta_r + 2a_1a_2\mathcal{X}\psi_r))((a_{21} - \alpha')\mathcal{X} + a_{22}\mathcal{Y} + (a_{23} + \beta'\cos\alpha)\mathcal{Z} + f(\xi_r + 2a_1a_2\mathcal{X}\phi_r)) \\
&+ a_{23}((a_{31} - \beta'\sin\alpha)\mathcal{X} + (a_{32} - \beta'\cos\alpha)\mathcal{Y} + a_{33}\mathcal{Z}) \\
&+ \alpha'(a_{11}\mathcal{X} + a_{12}\mathcal{Y} + a_{13}\mathcal{Z} + a_1f(\theta_r + a_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1a_2\mathcal{X}^2\psi_r)) \\
&- (a_{31}\mathcal{X} + a_{32}\mathcal{Y} + a_{33}\mathcal{Z})\beta'\cos\alpha - (-1)^r\nu a_1^2f(\xi_r + 2a_1a_2\mathcal{X}\phi_r), \tag{4.150}
\end{aligned}$$

$$\begin{aligned}
G_2 &= (b_{32} - b_{23})(b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}) \\
&- (b_{21} - b_{12} + a_1g(-a_1\varepsilon_r + b_2\phi_r - a_1b_2\mathcal{Y}\psi_r + (-1)^r a_1\mathcal{X}\xi_r + (-1)^r a_1^2b_2\mathcal{X}^2\phi_r)) \\
&\times (b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z} + a_1g(\theta_r + b_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1b_2\mathcal{X}^2\psi_r)), \tag{4.151}
\end{aligned}$$

$$\begin{aligned}
R_3 &= a'_{31}\mathcal{X} + a_{32}\mathcal{Y} + a'_{33}\mathcal{Z} + a_{31}(a_{11}\mathcal{X} + (a_{12} + \alpha')\mathcal{Y} + (a_{13} + \beta'\sin\alpha)\mathcal{Z} \\
&+ a_1f(\theta_r + a_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1a_2\mathcal{X}^2\psi_r)) \\
&+ a_{32}((a_{21} - \alpha')\mathcal{X} + a_{22}\mathcal{Y} + (a_{23} + \beta'\cos\alpha)\mathcal{Z} + f(\xi_r + 2a_1a_2\mathcal{X}\phi_r)) \\
&+ a_{33}((a_{31} - \beta'\sin\alpha)\mathcal{X} + (a_{32} - \beta'\cos\alpha)\mathcal{Y} + a_{33}\mathcal{Z}) \\
&+ (a_{11}\mathcal{X} + a_{12}\mathcal{Y} + a_{13}\mathcal{Z} + a_1f(\theta_r + a_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1a_2\mathcal{X}^2\psi_r))\beta'\sin\alpha \\
&+ (a_{11}\mathcal{X} + a_{12}\mathcal{Y} + a_{13}\mathcal{Z} + f(\xi_r + 2a_1a_2\mathcal{X}\phi_r))\beta'\cos\alpha, \tag{4.152}
\end{aligned}$$

$$\begin{aligned}
G_3 &= (b_{13} - b_{31})(b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z} + a_1g(\theta_r + b_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1b_2\mathcal{X}^2\psi_r)) \\
&- (b_{32} - b_{23})(b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z} + g(\xi_r + 2a_1b_2\mathcal{X}\phi_r)), \tag{4.153}
\end{aligned}$$

$$\begin{aligned}
&b'_{11}\mathcal{X} + b'_{12}\mathcal{Y} + b'_{13}\mathcal{Z} + (a_1g)'(\theta_r + b_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1b_2\mathcal{X}^2\psi_r) \\
&+ a_1g(a'_1\mathcal{Y}\varepsilon_r + b'_2\mathcal{Y}\phi_r + a'_1b_2\mathcal{Y}^2\psi_r - (-1)^r a'_1\mathcal{X}\mathcal{Y}\xi_r - (a_1b_2)'\mathcal{X}\psi_r - (-1)^r a'_1a_1b_2\mathcal{X}^2\mathcal{Y}\phi_r) \\
&+ (\alpha'\mathcal{Y} + \mathcal{Z}\beta'\sin\alpha)(b_{11} - a_1g(\zeta_r + 2a_1b_2\mathcal{X}\psi_r)) \\
&+ ((a_{21} - \alpha')\mathcal{X} + a_{22}\mathcal{Y} + (a_{23} + \beta'\cos\alpha)\mathcal{Z} + f(\xi_r + 2a_1a_2\mathcal{X}\phi_r)) \\
&\times (b_{12} + a_1g(a_1\varepsilon_r + b_2\phi_r + a_1b_2\mathcal{Y}\psi_r - (-1)^r a_1(\mathcal{X}\xi_r + a_1b_2\mathcal{X}^2\phi_r))) \\
&+ b_{13}((a_{31} - \beta'\sin\alpha)\mathcal{X} + (a_{32} - \beta'\cos\alpha)\mathcal{Y} + a_{23}\mathcal{Z}) \\
&+ (a_{22} + a_{33} + f(a_1\zeta_r + 2a_1^2a_2\mathcal{X}\psi_r)) \\
&\times (b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z} + a_1g(\theta_r + b_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1b_2\mathcal{X}^2\psi_r)) \\
&- (\alpha' + a_{12} + a_1f(a_1\varepsilon_r + a_2\phi_r + a_1a_2\mathcal{Y}\psi_r - (-1)^r a_1(\mathcal{X}\xi_r + a_1a_2\mathcal{X}^2\phi_r))) \\
&\times (b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z} + g(\xi_r + 2a_1b_2\mathcal{X}\phi_r)) - (a_{13} + \beta'\sin\alpha)(b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}) \\
&- (a_{11}\mathcal{X} + a_{12}\mathcal{Y} + a_{13}\mathcal{Z} + a_1f(\theta_r + a_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1a_2\mathcal{X}^2\psi_r)) \\
&\times (b_{22} + b_{33} + g(a_1\zeta_r + 2a_1^2b_2\mathcal{X}\psi_r)) \\
&- (-1)^r\eta a_1^3g(\theta_r + b_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1b_2\mathcal{X}^2\psi_r) = 0, \tag{4.154}
\end{aligned}$$

$$\begin{aligned}
& b'_{21}\mathcal{X} + b'_{22}\mathcal{Y} + b'_{23}\mathcal{Z} + g'(\xi_r + 2a_1b_2\mathcal{X}\phi_r) + g(a'_1\mathcal{Y}\zeta_r + 2(a_1b_2)'\mathcal{X}\phi_r + 2a'_1a_1b_2\mathcal{X}\psi_r) \\
& + (b_{21} + 2a_1b_2g\phi_r)(a_{11}\mathcal{X} + (a_{12} + \alpha')\mathcal{Y} + (a_{13} + \beta'\sin\alpha)\mathcal{Z}) \\
& + a_1f(\theta_r + a_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1a_2\mathcal{X}^2\psi_r) + (-\alpha'\mathcal{X} + \mathcal{Z}\beta'\cos\alpha)(b_{22} + a_1g(\zeta_r + 2a_1b_2\mathcal{X}\psi_r)) \\
& + b_{23}((a_{31} - \beta'\sin\alpha)\mathcal{X} + (a_{32} - \beta'\cos\alpha)\mathcal{Y} + a_{33}\mathcal{Z}) \\
& + (\alpha' - a_{21} - 2a_1a_2f\phi_r)(b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z} + a_1g(\theta_r + b_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1b_2\mathcal{X}^2\psi_r)) \\
& - (a_{22} + a_1f(\zeta_r + 2a_1a_2\mathcal{X}\psi_r))(b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z} + g(\xi_r + 2a_1b_2\mathcal{X}\phi_r)) \\
& - (a_{23} + \beta'\cos\alpha)(b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}) \\
& + (b_{22} + a_1g(\zeta_r + 2a_1b_2\mathcal{X}\psi_r))(a_{21}\mathcal{X} + a_{22}\mathcal{Y} + a_{23}\mathcal{Z} + f(\xi_r + 2a_1b_2\mathcal{X}\phi_r)) \\
& - (-1)^r\eta a_1^2g(\xi_r + 2a_1b_2\mathcal{X}\phi_r) = 0
\end{aligned} \tag{4.155}$$

and

$$\begin{aligned}
& b'_{31}\mathcal{X} + b'_{32}\mathcal{Y} + b'_{33}\mathcal{Z} + b_{31}(a_{11}\mathcal{X} + (a_{12} + \alpha')\mathcal{Y} + (a_{23} + \beta'\sin\alpha)\mathcal{Z}) \\
& + a_1f(\theta_r + a_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1a_2\mathcal{X}^2\psi_r) \\
& + b_{32}((a_{21} - \alpha')\mathcal{X} + a_{22}\mathcal{Y} + (a_{23} + \beta'\cos\alpha)\mathcal{Z} + f(\xi_r + 2a_1b_2\mathcal{X}\phi_r)) \\
& + (\beta'\sin\alpha - a_{31})(b_{11}\mathcal{X} + b_{12}\mathcal{Y} + b_{13}\mathcal{Z} + a_1g(\theta_r + b_2\mathcal{Y}\phi_r - \mathcal{X}\zeta_r - a_1b_2\mathcal{X}^2\psi_r)) \\
& + (\beta'\cos\alpha - a_{32})(b_{21}\mathcal{X} + b_{22}\mathcal{Y} + b_{23}\mathcal{Z} + g(\xi_r + 2a_1b_2\mathcal{X}\phi_r)) - a_{33}(b_{31}\mathcal{X} + b_{32}\mathcal{Y} + b_{33}\mathcal{Z}) \\
& + b_{33}(a_{31}\mathcal{X} + a_{32}\mathcal{Y} + a_{33}\mathcal{Z}) - b_{33}(\mathcal{X}\beta'\sin\alpha + \mathcal{Y}\beta'\cos\alpha) = 0.
\end{aligned} \tag{4.156}$$

From the coefficients of $\mathcal{X}^2\phi_r$ in $\Phi_{1\mathcal{Z}} = \Phi_{3\mathcal{X}}$, we get

$$(a_{23} + \beta'\cos\alpha)a_2f + \mu_0b_{23}b_2g = 0. \tag{4.157}$$

While, from the coefficients of $\mathcal{X}^2\phi_r$ in $\Psi_1 = 0$, we have

$$-(a_{23} + \beta'\cos\alpha)b_2g + a_2fb_{23} = 0. \tag{4.158}$$

Since the functions a_2 , b_2 , f and g are nonzero, we obtain that

$$\begin{cases} a_{23} = -\beta'\cos\alpha, \\ b_{23} = 0. \end{cases} \tag{4.159}$$

The coefficients of ϕ_r in $\Phi_{1\mathcal{Z}} = \Phi_{3\mathcal{X}}$ suggest that

$$a_{32} = -\beta'\cos\alpha. \tag{4.160}$$

The coefficients of ζ_r in $\Phi_{2\mathcal{Z}} = \Phi_{3\mathcal{Y}}$ say that

$$b_{32} = 0. \tag{4.161}$$

Thus the coefficients of ξ_r in $\Psi_3 = 0$ imply

$$\beta' = 0. \tag{4.162}$$

Substituting (4.159)-(4.162) into (4.148)-(4.156), we can simplify the calculation. One shows that

$$\begin{cases} a_{13} = a_{31} = 0, \\ b_{ij} = 0 \quad \text{for } i, j = 1, 2, 3, \end{cases} \tag{4.163}$$

$$\begin{cases} a_{21} = -a_{12} = \alpha' = k, \\ a_{11} = a_{22} = a_{33} = 0, \end{cases} \tag{4.164}$$

and

$$\begin{cases} f = c_1e^{(-1)^r\nu a_1^2t}, \\ g = c_2e^{(-1)^r\eta a_1^2t}, \\ b_2 = a_2 = \text{constant}, \end{cases} \tag{4.165}$$

where k , c_1 and c_2 are constants. Thus one easily shows that

$$\begin{aligned}
p = & \rho \{ -a_1^2 e^{(-1)^r 2\nu a_1^2 t} \left[\frac{1}{2} (4b\delta_{0,r} + c\delta_{1,r}) c \mathcal{X}^2 + a_2 (2\delta_{0,r}(ab+c) + \delta_{1,r} c \cos b) \mathcal{X}^3 \right. \\
& + a_1^2 a_2^2 (4a\delta_{0,r} + \delta_{1,r}) \frac{\mathcal{X}^4}{2} + a_2 (2\delta_{0,r}(ab-c) + \delta_{1,r} c \sin b) \mathcal{X} \mathcal{Y} + (\delta_{1,r} c c_1 \sin(b-b_1) \\
& + 2\delta_{0,r}(bc_1 - b_1 c)) \mathcal{X} + \frac{a_1 a_2}{2} (2\delta_{0,r}(c_1 - ab_1) - \delta_{1,r} c_1 \sin b_1) \mathcal{X}^2 \Big] + \frac{k^2}{2} \mathcal{X}^2 \\
& - e^{(-1)^r \nu a_1^2 t} (a_1 a_2 \phi - 2k) (\mathcal{X} \xi + a_1 a_2 e^{(-1)^r \nu a_1^2 t} \mathcal{X}^2 \phi_r) - (-1)^r e^{(-1)^r 2\nu a_1^2 t} [a_e^{(-1)^r 2\nu a_1^2 t} \\
& \times a_2 e^{(-1)^r \nu a_1^2 t} (a_1 a_2 \mathcal{Y} \phi_r \psi_r - \frac{a_2}{2} (a_1 (4a\delta_{0,r} + \delta_{1,r}) \mathcal{Y}^2 + \phi_r^2) + \theta_r \zeta_r \\
& - (\delta_{0,r} 2(ab_1 + c_1) + \delta_{1,r} c_1) \mathcal{Y}) + k(\varepsilon_r + a_2 \mathcal{Y} \psi_r - \frac{a_2}{a_1} \phi_r)] - \frac{1}{2} e^{(-1)^r 2\nu a_1^2 t} \xi_r^2 + \frac{k^2}{2} \mathcal{Y}^2 \Big\} \\
& + \mu_0 \rho \{ a_1 c_2 e^{(-1)^r \eta a_1^2 t} (-a_1 \xi_r \varepsilon_r + a_2 \xi_r \phi_r - a_1 a_2 \mathcal{Y} \xi_r \psi_r) \mathcal{X} \\
& + ((-1)^r a_1 \xi_r^2 - 2a_1^2 a_2 \phi_r \varepsilon_r + 2a_1 a_2^2 \phi_r^2 - 2a_1^2 a_2^2 \mathcal{Y} \phi_r \psi_r) \frac{\mathcal{X}^2}{2} \\
& + (-1)^r a_1^2 a_2 \mathcal{X}^3 \xi_r \phi_r + (-1)^r a_1^3 a_2^2 \frac{\mathcal{X}^4}{2} \phi_r^2 - \frac{a_1}{2} c_2 e^{(-1)^r 2\eta a_1^2 t} (\theta_r + a_2 \mathcal{Y} \phi_r)^2 - 2a_1^2 a_2 c_2 e^{(-1)^r 2\eta a_1^2 t} \\
& \times (\delta_{0,r} [\frac{1}{2a_1} (b_1 e^{2a_1 \mathcal{Y}} - a_1 c_1 e^{-2a_1 \mathcal{Y}}) - (c_1 + ab_1) \mathcal{Y}] + \delta_{1,r} [\frac{c_1}{2a_1} \sin b_1 \sin^2(a_1 \mathcal{Y}) \\
& + \frac{c_1}{b_1} \cos b_1 (\frac{a_1 \mathcal{Y}}{2} + \frac{\sin(2a_1 \mathcal{Y})}{4})]) \\
& - 2a_2^2 c_2 e^{(-1)^r 2\eta a_1^2 t} (\delta_{0,r} [-a_1^2 \mathcal{Y}^2 + \frac{a_1 \mathcal{Y}}{2} (e^{2a_1 \mathcal{Y}} - e^{-2a_1 \mathcal{Y}}) - \frac{1}{4} (e^{2a_1 \mathcal{Y}} + e^{-2a_1 \mathcal{Y}})] \\
& - \delta_{1,r} [\frac{5}{8} \sin(2a_1 \mathcal{Y}) - \frac{\sin^2(a_1 \mathcal{Y})}{2} - \frac{a_1 \mathcal{Y} \sin^2(a_1 \mathcal{Y})}{2} + \frac{a_1 \mathcal{Y}}{4}]) \Big\} \tag{4.166}
\end{aligned}$$

modulo the transformation in (1.9).

Theorem 4.6. *Let k , l , a_1 , a_2 , c_1 and c_2 be any real constants. Then we have the following solutions of the MHD equations (1.1)-(1.4):*

$$\begin{cases} u = k \mathcal{X} \sin(kt + l) - k \mathcal{Y} \cos(kt + l) - c_1 e^{(-1)^r \nu a_1^2 t} (\xi_r + 2a_1 a_2 \mathcal{X} \phi_r) \sin(kt + l) \\ \quad + a_1 c_1 e^{(-1)^r \nu a_1^2 t} (\theta_r + a_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 a_2 \mathcal{X}^2 \psi_r) \cos(kt + l), \\ v = k \mathcal{X} \cos(kt + l) - k \mathcal{Y} \sin(kt + l) + a_1 c_1 e^{(-1)^r \nu a_1^2 t} (\theta_r + a_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 a_2 \mathcal{X}^2 \psi_r) \\ \quad \times \sin(kt + l) + c_1 e^{(-1)^r \nu a_1^2 t} (\xi_r + 2a_1 a_2 \mathcal{X} \phi_r) \cos(kt + l) \\ w = 0, \end{cases} \tag{4.167}$$

$$\begin{cases} H^1 = a_1 c_2 e^{(-1)^r \eta a_1^2 t} (\theta_r + a_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 a_2 \mathcal{X}^2 \psi_r) \cos(kt + l) \\ \quad - c_2 e^{(-1)^r \eta a_1^2 t} (\xi_r + 2a_1 a_2 \mathcal{X} \phi_r) \sin(kt + l), \\ H^2 = a_1 c_2 e^{(-1)^r \eta a_1^2 t} (\theta_r + a_2 \mathcal{Y} \phi_r - \mathcal{X} \zeta_r - a_1 a_2 \mathcal{X}^2 \psi_r) \sin(kt + l) \\ \quad + c_2 e^{(-1)^r \eta a_1^2 t} (\xi_r + 2a_1 a_2 \mathcal{X} \phi_r) \cos(kt + l), \\ H^3 = 0 \end{cases} \tag{4.168}$$

and p is given by (4.166).

Acknowledgement: I would like to thank Professor Xiaoping Xu for his advice and suggesting this research topic.

References

- [1] J. F. Gerbeau, C. Le Bris and T. Lelievre, *Mathematical Methods for the Magnetohydrodynamics of Liquid Crystals*, Oxford Science Publication, 2006.
- [2] M. Cannone, Q.L. Chen, and C. Miao, A losing estimates for the ideal MHD equations with application to blow-up criterion, *SIAM J. Math. Anal.* **38** (2007), 1847–1859.
- [3] M. Cannone, C. Miao, N. Prioux, and B.Q. Yuan, *The Cauchy Problem for the Magneto-Hydrodynamic System, Self-Similar Solutions of Nonlinear PDE*, Vol. **74**, Banach Center Publications, Institute of Mathematics Polish Academy of Sciences, Warszawa, 2006.
- [4] Q. Chen, C. Miao, and Z. Zhang, The Beale-Kato-Majda criterion for the 3D magnetohydrodynamics equations, *Comm. Math. Phys.* **275** (2007), 861–872.
- [5] J. Wu, Lower bounds for an integral involving fractional Laplacians and the generalized Navier- Stokes equations in Besov spaces, *Comm. Math. Phys.* **263** (2006), 803–831.
- [6] J. Wu, Regularity criteria for the generalized MHD equations, *Comm. Partial Diff. Equat.* **33** (2008), 285–306.
- [7] J. Zhang, Wellposedness for the magnetohydrodynamics equation in critical space, *Appl. Anal.* **87** (2008), no. 7, 773–785.
- [8] M. Brio and C. C. Wu, An upwind differencing scheme for the equations of ideal magnetohydrodynamics, *J. Comput. Phys.* **75**(1988), 400–422 .
- [9] S. Zachary and P. Colella, A higher-order Godunov method for the equations of ideal magnetohydrodynamics, *J. Comput. Phys.* **99** (1992), 341–347.
- [10] P. L. Roe and D. S. Balsara, Notes on the eigensystem of magnetohydrodynamics, *SIAM J. Appl. Math.* **56** (1996), 57–67.
- [11] W. Dai, P. R. Woodward, An approximate Riemann solver for ideal magnetohydrodynamics, *J. Comput. Phys.* **111** (1994), 354–372.
- [12] J. Croisille, R. Khanfir, and G. Chanteur, Numerical simulation of the MHD equations by a kinetic-type method, *J. Sci. Comput.* **10** (1995), 81–92.
- [13] T. Linde, *A Three-Dimensional Adaptive Multifluid MHD Model of the Heliosphere*, Ph.D. thesis, University of Michigan, 1998.
- [14] R. Myong, *Theoretical and Computational Investigation of Nonlinear Waves in Magnetohydrodynamics*, Ph.D. thesis, University of Michigan, 1996.
- [15] G. Tóth and D. Odstreil, Comparison of some flux corrected transport and total variation diminishing numerical schemes for hydrodynamic and magnetohydrodynamic problems, *J. Comput. Phys.* **128** (1996), 82–112.
- [16] Kenneth G. Powell, Philip L. Roe, Timur J. Linde, Tamas I. Gombosi, and Darren L. De Zeeuw, A solution-adaptive upwind scheme for ideal magnetohydrodynamics, *J. Comput. Phys.* **154** (1999), 284–309.
- [17] Dongwook Lee, Anil E. Deane, An unsplit staggered mesh scheme for multidimensional magnetohydrodynamics, *J. Comput. Phys.* **228** (2009), 952–975.
- [18] D.S. Ryu, T.W. Jones, Numerical magnetohydrodynamics in astrophysics: algorithm and tests for one-dimensional flow, *Astrophys. J.* **442** (1995), 228–253.

- [19] M.D. Gunzburger, A.J. Meir and J.S. Peterson, On the existence and uniqueness and finite element approximation of solutions of the equations of stationary incompressible magnetohydrodynamics, *Math. Comp.* **56** (1991), 523–563.
- [20] U. Hasler, A. Schneebeli and D. Schötzau, Mixed finite element approximation of incompressible MHD problems based on weighted regularization, *Appl. Numer. Math.* **51** (2004), 19–45.
- [21] D. Schötzau, Mixed finite element methods for stationary incompressible magnetohydrodynamics, *Numer. Math.* **96** (2004), 771–800.
- [22] A. Prohl, Convergent finite element discretizations of the nonstationary incompressible magnetohydrodynamics system, *ESAIM: M2AN* **42** (2008), 1065–1087.
- [23] K. S. Kang, and D. E. Keyes, Implicit symmetrized streamfunction formulations of magnetohydrodynamics, *Int. J. Numer. Meth. Fluids* **58** (2008), 1201–1222.
- [24] Bhimsen K. Shivamoggi, Impulse formulations of Hall magnetohydrodynamic equations, *Phys. Lett. A* **373** (2009), 708–710.
- [25] Rostislav Vodák, Local solvability of equations of magnetohydrodynamics on the whole space, *Nonl. Analysis: Real World Appl.* **10** (2009), 1424–1442.
- [26] T. G. Cowling, Magnetohydrodynamics, *Rep. Prog. Phys.* **25**(1962), 244–286.
- [27] N. H. Ibragimov, *Lie Group Analysis of Differential Equations*, Vol 2, CRC Handbook, CRC Press, 1995.
- [28] M. C. Nucci, Group analysis for MHD equations, in: *Atti Sem. Mat. Fis. Univ. Modena*, 1984, V. XXXIII, 21–34.
- [29] V. O. Popovych, On Lie reduction of the MHD equations to ordinary differential equations, *Proceedings of the Second International Conference "Symmetry in Nonlinear Mathematical Physics" (Kyiv, July 7–13, 1997)*, Kyiv, Institute of Mathematics, 1997, Vol. 1, 227–231.
- [30] V. O. Popovych, On classes of Lie solutions of MHD equations, expressed via the general solution of the heat equations, *J. Nonl. Math. Phys.* **4** (1997), 149–151.
- [31] V. O. Popovych, Lie submodels of rank 1 for MHD equations, *Proc. Inst. Math. NAS Ukraine* **30** (2000), 190–195.
- [32] X. Xu, Asymmetric and moving-frame approaches to Navier-Stokes equations, *Quart. Appl. Math.* **67** (2009), 163–193.